

1 **Runoff, nutrients, sediment and salt yields in an irrigated**
2 **watershed in southern Navarre (Spain)**

3
4 *D. Merchán^a, J. Casal^a, J. Del Valle de Lersund^b, M.A. Campo-Bescós^a, R. Giménez^a,*
5 *B. Preciado^c, A. Lafarga^c*

6
7 ^a Department of Projects and Rural Engineering – Public University of Navarre; Edificio Los
8 Olivos - Campus de Arrosadía, 31006 Pamplona, Navarra, Spain

9 ^b Department of Rural Development, Environment and Local Administration – Government of
10 Navarre, 31005 Pamplona, Navarra, Spain

11 ^c Navarre Institute of Agricultural and Food Technologies and Infrastructures, Edificio Peritos -
12 Avda. Serapio Huici 22. 31610 Villava, Navarra, Spain

13
14
15 **ABSTRACT**

16
17 The environmental impact of irrigated agriculture on water quality was assessed in
18 Landazuria watershed (Navarre, northeast Spain), a 479.5 ha watershed with 53% of
19 irrigated agricultural land. In the framework of a long-term monitoring program,
20 precipitation and discharge were measured at 10-minutes intervals and compound daily
21 water samples were collected during the agricultural years (September to August) 2007-
22 2016, and analysed for nitrate (NO_3^-), phosphate (PO_4^{3-}), sediment and total dissolved
23 solids (TDS) concentrations. Typical agricultural management (including crop surfaces,
24 irrigation and fertilization rates) was obtained from inquiries to farmers. Concentration
25 and yield of the studied variables presented a high degree of variation, both intra- and
26 inter-annual. Median concentration for the entire study period were 185, <0.05, 31 and
27 2284 mg L⁻¹ for NO_3^- , PO_4^{3-} , sediment and TDS, respectively. NO_3^- -N and PO_4^{3-} -P yields
28 averaged 74 and 0.04 kg ha⁻¹ year⁻¹, respectively. NO_3^- -N yield was higher than in other

29 agricultural land uses in Navarre and in the order of magnitude of other irrigated areas
30 in the Middle Ebro Valley. $\text{PO}_4^{3-}\text{-P}$ yield was in the same order of magnitude than in
31 rainfed watersheds in Navarre but lower than in intensively grazed watersheds. Sediment
32 yield was extremely variable, averaging $360 \text{ kg ha}^{-1} \text{ year}^{-1}$, with 44 % of the total
33 measured load recorded in a few days. It was in the lower range of those measured in
34 Navarre for rainfed agriculture and similar to those estimated in other irrigated areas of
35 the Middle Ebro River. TDS concentration presented a significant decreasing trend since
36 available salts were being washed out, while TDS yield averaged $1.8 \text{ Mg ha}^{-1} \text{ year}^{-1}$.
37 Long-term monitoring of irrigated areas is required to understand pollution processes in
38 these agroecosystems and to adequately characterize the environmental impact of
39 current agricultural practices on water quality, in order to implement, and adequately
40 assess, measures to reduce agricultural pollution.

41

42 **Keywords:** watershed, agricultural pollution, nitrate, phosphate, soil loss, TDS.

43

44

45

46

47 **1. Introduction**

48 More than 1.5 billion ha (about 12% of the world's land area) are used for crop
49 production. Rainfed agriculture is the predominant agricultural production system, but
50 increasing climate variability is bringing greater uncertainty in the production levels.
51 Current productivity in rainfed systems is, on average, little more than half of its potential
52 (FAO, 2013). However, agricultural production has grown between 2 and 4% per year
53 over the last 50 years, while the cultivated area has grown by only 1% annually. More
54 than 40% of the increase in food production has come from irrigated areas.

55

56 Irrigation has many advantages over rainfed agriculture, such as increased productivity,
57 higher diversity of crops, more reliable harvests, or regional economic security (e.g.,
58 Duncan et al., 2008). For these reasons, a global increase in irrigated surface has been
59 observed, especially in developing countries, where it doubled between 1962 and 1998
60 (FAO, 2003a). In Spain, the increase of irrigated area has been moderate but significant,
61 with 7% more irrigated land between 1990 and 2009 according to the Spanish Ministry
62 of Agriculture and Fisheries, Food and Environment (MAPAMA, 2017). In fact, irrigated
63 agriculture has been a key factor in the agrarian Spanish system as it provides more
64 than 50% of the final agrarian production with only 13% of the surface. According to
65 official estimates, in Spain an irrigated hectare produces, on average, six times more
66 than a rainfed hectare, and generates four times more income. In Navarre (10,391 km²,
67 northern Spain), irrigated surface has increased in recent years to over 110,000 ha
68 (approximately 25% growth between 2000 and 2015; DDRMAAL, 2017), being
69 pressurized districts those implemented in the new irrigated land.

70

71 There is no question about the value of irrigated agriculture but there is an increasing
72 trend to make it accountable for its impacts on the environment (e.g., Stockle, 2001).
73 Agricultural land use is regarded as the main source of diffuse pollution (Novotny, 1999),
74 and it has a wide range of associated environmental impacts such as changes in

75 landscapes and plant and animal communities, and the deterioration of soil, water and
76 air quality (Stoate et al., 2001; Merrington et al., 2002). Specifically, irrigated agriculture
77 imposes severe pressure on the environment, as it accounts for the consumption of 70%
78 of global water resources (FAO, 2003b), being the main reason behind the construction
79 of most dams or aquifers overexploitation. Apart from the effects on the withdrawal water
80 body, irrigation return flows can also cause hydrological changes in the receiving ones.
81 Specific environmental problems in waters downstream of the irrigated areas are related,
82 among others, to nutrients, sediments or salts.

83

84 Nitrate pollution is a major concern in irrigated areas since high nitrate concentrations
85 have long been regarded as a threat for human health and ecosystems (e.g., Sutton et
86 al., 2011). Despite the fact that nitrate leaching varies considerably with climatic
87 conditions (e.g., Elmi et al., 2004), the actual impact of N pollution depends on specific
88 features of the area such as the soil types (Kyllmar et al., 2014), the presence of reducing
89 conditions in aquifers (Rivett et al., 2008) and the irrigation/fertilization management
90 (Quemada et al., 2013). On the other hand, the mobility of phosphorus is rather limited,
91 especially in arid or semi-arid regions (Brady and Weil, 2008). In these soils, only a small
92 fraction of P is in soluble reactive form (phosphate) and can be incorporated into plants.
93 For this reason, P losses are normally related to soil or sediment losses (Edwards and
94 Withers, 1998). As a consequence, P pollution is greatly conditioned by erosion,
95 especially in arid and semi-arid areas.

96

97 Erosion processes in agricultural land tend to be significantly higher than in other land
98 uses (García-Ruiz et al., 2015). Erosion removes preferentially the fine fraction of soils,
99 which is enriched in nutrients and organic matter (Merrington et al., 2002). Soil erosion
100 rates are extremely variable (García-Ruiz et al., 2015) and depend on both natural
101 factors (climate, slope, soils, bedrock...) and agronomic management (cover, tillage...).
102 Finally, the leaching of salts is a requirement of irrigated agriculture (Letey et al., 2011)

103 since its build-up in soils can be deleterious for plants, decrease productivity and even
104 force the abandonment of cultivation. However, leached salts will reach water bodies
105 downstream, affecting its quality for human consumption or ecosystem uses (Nielsen et
106 al., 2003). The amount of salts leached depends on different factors such as climate,
107 hydrogeological conditions or irrigation management (Merchán et al., 2015c).

108

109 In Navarre, the environmental impact of agriculture is investigated in a network of
110 experimental watersheds (Fig. 1) implemented by the former *Department of Agriculture,*
111 *Livestock and Food* of the Government of Navarre. This network includes representative
112 agricultural land-uses in the region (Casalí et al., 2008, 2010). An irrigated watershed
113 (Landazuria) was included in the monitoring network in 2006. The climatic, geologic and
114 agronomic characteristics of this watershed make it representative of the recent
115 pressurized irrigated areas in the Middle Ebro Valley, with over 900,000 ha dedicated to
116 irrigated agriculture, and approximately half of this surface being pressurized irrigation
117 systems (CHE, 2017). In addition, in the framework of the project LIFE-Nitrates (LIFE+10
118 ENV/ES/478), a consortium of public institutions constituted by the Government of
119 Navarre (GN), Environmental Management of Navarre (GAN), Navarre Institute of
120 Agricultural and Food Technologies and Infrastructures (INTIA) and CRANA Foundation
121 conducted a detailed study on the “Impacts of agricultural practices on nitrate pollution
122 of continental waters” (www.life-nitratos.eu/). One of the study sites investigated in this
123 project was the irrigated watershed monitored by the Government of Navarre.

124

125 In this paper we present the data obtained during the agricultural years 2007-2016 in the
126 irrigated watershed (Landazuria). Given the high variability in climatic and agronomic
127 conditions in specific study cases and the different processes affecting different
128 pollutants, the analysis of detailed and long-term temporal series of a wide set of
129 variables is paramount to better understand the pollution of water bodies as a
130 consequence of agricultural land use, particularly in irrigated areas. The main objectives

131 were: (1) to estimate the effects of irrigated agriculture on water quality, specifically in
132 terms of nitrate, phosphate, sediment and salts concentration in the watershed outlet
133 and exported yields; (2) to determine the controlling factors explaining that behaviour;
134 and (3) to contextualize the obtained estimations and inferred controlling factors with
135 those reported in other irrigated and rainfed watersheds, paying especial attention to the
136 difference between them.

137 **2. Methods**

138 *2.1. Experimental watershed*

139 Landazuria watershed covers an area of 479.5 ha and is located in southern Navarre
140 (Fig. 1). It is relatively flat, with slopes between 3.5 and 5%. A single 1st order stream
141 drains the watershed. The geographical coordinates of the watershed outlet are
142 42°15'3.5"N and 1°35'3.4"W. According to data collected for the period 1992-2016
143 around 5 km south (meteorological station Bardenas-El Yugo, Government of Navarra)
144 the climate in the study zone is Dry Mediterranean. Average annual temperature is 14
145 °C, but it can reach values as low as -8 °C in winter and as high as 41 °C in summer.
146 Annual precipitation is 426 ± 114 mm (average ± standard deviation) whereas annual
147 reference evapotranspiration (FAO Penman-Monteith, Allen et al., 1998) is 1369 ± 101
148 mm, i.e., more than three times higher and much less variable.

149

150 The geology in Landazuria is represented by Tertiary and Quaternary materials. The
151 Tertiary materials appear as a bottom layer several hundred metres thick, composed of
152 alternating gypsum, and red clays, with occasional intercalations of fine (centimetres to
153 decimetres) limestone layers (DOPTC, 2003a, 2003b). The Quaternary materials cover
154 in most of the watershed surface the Tertiary materials, and they are composed mainly
155 by detrital sediments, gravels with some limestone clasts, alternating with sands, silt and
156 clays (glacis) of Pleistocene-Holocene age. The synclinal structure and extremely low
157 hydraulic conductivity of the Tertiary materials ($<10^{-8}$ m s⁻¹; DOPTC, 2003a, 2003b)
158 avoids deep percolation of water within the watershed.

159

160 According to a detailed survey including 78 direct observations (Government of Navarre,
161 2005), soils developed in Landazuria present mainly clay loam or silt loam textures and
162 are deep, with the exception of eroded hills (Fig. 1c). The most common series
163 correspond to Typic Haplustepts and Typic Calciustolls, although other series do appear
164 (Fig. 2 and Table 1). Organic matter content ranged between 1.7 and 2.7 % while pH
165 ranged between 8.1 and 8.9. Landazuria soils presented in general low salinity, with
166 some slight to moderate salinity level in the valley bottoms and in soils developed over
167 Tertiary marls.

168

169 Around 88.3 % of Landazuria is cultivated land and 11.7 % streams, riparian vegetation,
170 bare soils, ways and rock outcrops (Fig. 1b). In the year 1999, a surface of 252.9 ha was
171 equipped for pressurized irrigation (59.7 % of cultivated land) whereas the rest of the
172 surface remained as rainfed agriculture (170.5 ha or 40.3 % of cultivated land). According
173 to face-to-face farmers' inquiries (INTIA, 2017), the main crops under irrigation where
174 maize, winter cereal, tomatoes and onions (Table 1). Rainfed surface was dominated by
175 barley, although it is worthy to mention that rainfed agriculture in Landazuria followed a
176 cultivation system in which the land is left bare one out of two years (Table 1).

177

178 The irrigation system more widely used in Landazuria is solid-set sprinkler irrigation
179 (89%), although minor surfaces of drip irrigation do exist (e.g., tomatoes). The pressure
180 is provided by the difference in elevation between a reservoir or several distribution
181 ponds and the agricultural plots. The origin of irrigation water is a neighbour watershed,
182 so irrigation implies in an additional input in Landazuria. Irrigation volumes applied were
183 estimated according to the recommendations provided by INTIA, based on the irrigation
184 requirement of the crops. To this end, reference evapotranspiration (ET_0) computed by
185 the FAO Penman-Monteith methodology (Allen et al., 1998) was used in combination
186 with crop coefficients to obtain crop evapotranspiration ($ET_C = K_C \cdot ET_0$). The amount of

187 water provided by precipitation was considered and an irrigation efficiency of 85% and
188 95% was assumed for sprinkler and drip irrigation systems, typical values used by the
189 agricultural extension service in the region.

190

191 Fertilization practices varied widely among crops and, especially, between irrigated and
192 rainfed surfaces. Detailed information about fertilization practices for the period 2007 –
193 2013 was available for this study (Table 1). According to the aforementioned farmers
194 inquiries (INTIA, 2017), nitrogen fertilization rates in the rainfed surface averaged 32 kg
195 N ha⁻¹ year⁻¹, while in the irrigated surface they averaged 208 kg N ha⁻¹ year⁻¹, that is,
196 6.5 times higher (Table 1). This fact combined with the fallow land proportion in the
197 rainfed surface and the existence of double cropping patterns in irrigated surface (Table
198 1) implies that rainfed crops fertilization accounted only for 4.3 % of N fertilization in the
199 study area.

200

201 Maize was the crop with highest fertilization rates, between 224 and 423 kg N ha⁻¹ year⁻¹
202 ¹ (average of 285 kg N ha⁻¹ year⁻¹). Maize fertilization was carried in two steps in most of
203 the cases, complex fertilizer as basal application (normally 18-46-0, 9-23-30 or 8-15-15)
204 followed by urea or liquid fertilizers (e.g., N32 solution) as topdressing. Irrigated winter
205 cereal (barley and wheat) received around 138 kg N ha⁻¹ year⁻¹ split in December and
206 February-March. Tomato received 218 kg N ha⁻¹ year⁻¹ in several applications, normally
207 complex fertilizers as basal application before seeding followed by two topdressing
208 applications of liquid N32 (fertigation). Onion received 168 kg N ha⁻¹ year⁻¹ normally in
209 two applications with significant contents of sulphur (e.g., ammonium nitrosulphate).
210 Among rainfed crops, barley received 48 kg N ha⁻¹ year⁻¹ normally in a unique basal
211 application as complex fertilizers. Fertilization rates regarding other nutrients such as P,
212 K or S were not systematically collected since they were out of the scope of the LIFE-
213 Nitrates project. Only partial information was available (for instance, that recorded when

214 complex NPK fertilizer were applied) and therefore it was not analysed in detail in this
215 study.

216

217 Considering all ploughing methods, for the period with available data, an average of 689
218 ha year⁻¹ were ploughed, i.e., 63% more than the arable land, since several plots were
219 ploughed in several occasions during the year (INTIA, 2017). 80.6 % of the total
220 ploughed surface was irrigated land, whereas 19.4 % was rainfed land, what implies that,
221 as an average, irrigated land was ploughed twice a year while rainfed land was not
222 ploughed each year, as expected given the high proportion of fallow land. This fact
223 together with the fertilization rates aforementioned denotes the intense use of irrigated
224 land in Landazuria in comparison with that of rainfed land. Ploughing actuations were
225 mainly carried out in two different periods: between February and April (47 % of ploughed
226 surface) and between August and October (22 %). The remaining months in the year
227 together accounted for 31 % of the ploughed surface.

228

229 Average productions for the period 2007-2013 were 11,850 kg ha⁻¹ for maize, over
230 90,000 kg ha⁻¹ for tomato, 68,500 kg ha⁻¹ for onion, 6,150 and 1,500 kg ha⁻¹ for irrigated
231 and rainfed wheat, respectively, and 5,600 and 2,100 kg ha⁻¹ for irrigated and rainfed
232 barley, respectively (INTIA, 2017). It is worthy to note the different productions attained
233 under irrigated and rainfed conditions given the local conditions in Landazuria.

234

235 Although a great degree of variation is expected depending on the specific study site,
236 due to its climatological, geological and agronomic characteristics, Landazuria
237 watershed may be regarded as representative of the new pressurized irrigation districts
238 that have been implemented in the Middle Ebro Valley during the last decades.

239

240 *2.2. Data collection*

241 The former *Department of Agriculture, Livestock and Food* of the Government of Navarre
242 installed a hydrological station at the watershed outlet in the summer of 2006. Since then,
243 water level was recorded at 10 minutes intervals. The discharge measurement device
244 consisted of an H-type flume. Discharge was calculated from water level data, which
245 were monitored using a pressure probe and data logger. According to the hydraulic
246 characteristics of the weir, the following rating curve was used to transform the water
247 level (h , m) in discharge (Q , L s⁻¹):

$$248 \quad Q = 449.11 \cdot h^3 + 622.19 \cdot h^2 + 8.4101 \cdot h + 0.2248 \quad [\text{Eq. 1}]$$

249 Water discharge was also directly measured for verification using a propeller-type current
250 meter and triangular and rectangular sharp-crested weirs. All measurement methods
251 yielded consistent results.

252

253 Water samples were taken every 6 hours from a hemispheric hollow, 0.66 m in diameter,
254 made in the downstream face of the H-type flume. For this purpose, an automatic
255 programmable sampler was used, consisting of 24 bottles (500 mL). The four samples
256 collected each day were mixed together prior to analysis to provide a representative daily
257 average sample for determining sediment and nutrient concentration (Isidoro et al.,
258 2003). Water samples were analysed following the standard methods for water quality
259 parameter at the Agricultural Laboratory of the Department of Agriculture and Food of
260 the Government of Navarre. Suspended sediment concentration and dissolved nitrate
261 (NO_3^-) and phosphate (PO_4^{3-}) concentrations were determined as well as the pH (Crison
262 52-02 probe) and the major dissolved constituents (Na^+ , K^+ , Ca^{2+} , Mg^{2+} , Cl^- , SO_4^{2-} , HCO_3^-
263 CO_3^{2-}). Cations were determined by inductively coupled plasma-optical emission
264 spectrometry (ICP-OES); Cl^- , SO_4^{2-} and NO_3^- by ionic chromatography technique; HCO_3^-
265 and CO_3^{2-} by acid-base volumetric technique; PO_4^{3-} by spectrophotometry (ammonium
266 molybdate); and suspended sediment by gravimetric technique (0.7 μm pore size). The
267 charge balance of the samples was determined and found to be within $\pm 10\%$ for most

268 of the samples, suggesting that all relevant constituents were considered. Total dissolved
269 solids (TDS) was then computed by addition of the individual dissolved constituents.

270

271 *2.3. Data treatment, statistical procedures and interpretation*

272 10-minute meteorological and discharge values were processed to hourly, daily and
273 monthly time-series for analysis. Daily discharge data and concentration was used to
274 estimate the daily load of nitrate-nitrogen (NO_3^- -N), phosphate-phosphorus (PO_4^{3-} -P),
275 sediment and salts. TDS is usually used to estimate salt content of a water sample.
276 However, in Landazuria TDS was only determined in a subset of the available samples
277 (1,721 out of 2,984). For this reason, the existing relationship between chloride and total
278 dissolved solids (Fig. 2) was used to generate a TDS value for every sample. This value
279 was used in the estimation of salt loads.

280 Several problems produced missing samples for specific days (e.g., equipment
281 malfunctioning, not enough flow for sample collection, etc.). In fact, for the 10-year study
282 period (3,653 days), a total of 2,984 water samples were collected (82 %). As a
283 consequence, in order to obtain monthly time-series, the following criteria was applied:

284 (1) To obtain monthly concentration values, the median was estimated with a 95%
285 confidence interval using all available samples for that specific month;

286 (2) For monthly load values:

287 (a) If more than 33% of the days in the month had available samples, an
288 estimation was computed assigning the median value for that month to the days
289 without sample.

290 (b) No estimation of the load was obtained for those months with less than 33%
291 of the days in the month with available samples.

292

293 For PO_4^{3-} and sediments, a significant proportion of the samples were below detection
294 limits (BDL), $<0.05 \text{ mg L}^{-1}$ for PO_4^{3-} and $<5 \text{ mg L}^{-1}$ for sediments. To avoid bias in the

295 estimation of concentration and yields, robust methods were used to deal with
296 concentrations BDL (Helsel and Hirsch, 2002).

297

298 As exposed previously, there were gaps in the time series due to different issues. For
299 this reason, a temporal resolution of weeks was selected to study the correlation between
300 the different variables. Out of the 10 year study period, 381 complete weeks were
301 sampled (73%). For each week, the precipitation, irrigation, and fertilization amounts
302 were computed or estimated. In addition, average concentration and total load exported
303 of studied variables were computed. A correlation matrix was obtained using the non-
304 parametric Spearman's rho test (Helsel and Hirsch, 2002).

305

306

307 **3. Results and discussion**

308 *3.1. Precipitation, reference evapotranspiration, irrigation and runoff*

309 During the study period (agricultural years 2007-2016), annual precipitation (P) ranged
310 from 309 mm (2012) to 631 mm (2013), whereas reference evapotranspiration (ET_0)
311 ranged from 1253 mm to 1558 mm in the years 2013 and 2009, respectively. Average
312 values were 437 ± 97 mm and 1380 ± 85 mm for P and ET_0 , respectively. There were
313 no significant differences with the long-term data (section 2.1). In addition, no significant
314 trends (Seasonal Kendall Test, Helsel and Hirsch, 2002) were detected for monthly or
315 annual data of P or ET_0 , neither in the study period nor in the long-term data. The lack
316 of long-term trends suggest that the dynamics observed were representative of the
317 natural variability in Landazuria.

318

319 P was, in some extent, evenly distributed throughout the year, with 29%, 25%, 29% and
320 17% of the annual value for autumn, winter, spring and summer, respectively (Table 2).
321 In fact, although summer presented the lowest precipitation, there were no significant
322 differences between seasons ($p > 0.05$). In contrast, ET_0 was not evenly distributed, with

323 12%, 12%, 34% and 42%, respectively. Estimated irrigation volumes required for crops
324 (I) averaged 337 mm year⁻¹, with 2%, 4%, 31% and 62% of the annual value for autumn,
325 winter, spring and summer, respectively. As a consequence of the irrigation volume
326 distribution, an even distribution of discharge volume (Q) throughout the year was
327 observed (Table 2, Fig. 3). In fact, Q tended to be higher during summer than during
328 winter, with averages of 29 ± 9 mm and 22 ± 9 mm, respectively. However, the
329 differences between summer and winter discharge were not significant (p>0.05).

330

331

332 During the irrigated season, a daily cycle in Landazuria discharge was detected (Fig. 4).
333 Discharge reached a maximum around 9:00 hours and a minimum around 21:00 hours.
334 This observation is related to the irrigation practice of farmers in Landazuria, who irrigate
335 mainly during the night since wind speed and relative humidity were more favourable for
336 sprinkler irrigation, minimizing evaporation and wind drift losses (Playán et al., 2005).
337 Similar observations have been reported in other pressurized irrigation districts in the
338 Middle Ebro Valley (e.g., Isidoro et al., 2003; Causapé et al., 2012).

339

340 Since the irrigation volumes required by crops were relatively constant during the years
341 covered by this study, the differences in the distribution of the daily discharge were
342 related to the precipitation regime of the different years (Table 3), i.e., the amount of
343 precipitation and its distribution throughout the year. For instance, the dry year 2012
344 presented the lowest median and quartiles whereas the humid year 2015 presented the
345 highest.

346

347 *3.2. Nitrate concentrations and nitrate-nitrogen yields*

348 Measured NO₃⁻ concentrations in Landazuria outlet were considerably high, over three
349 times the European guideline for continental waters (50 mg L⁻¹) and even higher (Fig. 8).
350 In several years, a sharp increase in NO₃⁻ concentration in early summer (May and June)

351 was observed coinciding with the high fertiliser applications in the irrigated plots. Similar
352 observations have been reported in other irrigated watersheds in the Middle Ebro Valley
353 (Isidoro et al., 2003; Merchán et al., 2013) where a crop with high N needs dominated
354 the irrigated surface (such as maize in the case of Landazuria).

355

356 Significant differences in NO_3^- concentration were observed between agricultural years
357 (Fig. 8). For instance, the year 2013 presented the highest median and quartiles. This
358 year combined high fertilization rates during the last years (Table 1) with a previous dry
359 year (2012, the driest year covered in the study period; Table 3). Indeed, NO_3^-
360 concentration tends to be higher after dry years (Burt et al., 2012). In contrast, the years
361 2015 and 2016 presented the lowest NO_3^- concentrations, being significantly lower than
362 those recorded in any other year ($p < 0.001$). Several explanation may justify this
363 observation. Firstly, humid years (such as 2013 and 2015 in Landazuria) may deplete
364 the NO_3^- pool available in soils or phreatic layers, contributing to lower concentrations in
365 following years. In addition, a subtle decrease in the surface of crops requiring high N
366 fertilisation (maize) is observed in the last years (Table 1), what may have promoted
367 increased NO_3^- extractions. Finally, most of the farmers in the study area received
368 training and attended informative sessions about Best Management Practices to reduce
369 NO_3^- leaching in the framework of the LIFE-Nitrates Project (INTIA and GAN, 2015a).
370 Probably, a combination of these explanations is behind the lower concentration in the
371 last two studied years. Unfortunately, there was not available information about the
372 possible shift of fertilisation practices to adequately assess the contribution of each
373 explanation.

374

375 Monthly nitrate-nitrogen (NO_3^- -N) loads varied widely intra- and inter-years, and in
376 general they were heavily conditioned by discharge (Table 3). For those years in which
377 a complete estimation was possible, an average NO_3^- -N yield of $39 \pm 13 \text{ kg N ha}^{-1} \text{ year}^{-1}$
378 ¹. However, N fertilisers applied to the irrigated surface accounted for 96 % of total

379 applications (see section 2.1). Thus, considering negligible the nitrogen leached from
380 rainfed and bare soils, yield averaged ca. $70 \pm 24 \text{ kg N ha}^{-1} \text{ year}^{-1}$ for the irrigated surface.
381 This figure implies that, as an average, a mass of about 35 % of applied N in the irrigated
382 area leaves the watershed (Table 3). The yield reached up to $114 \text{ kg NO}_3^- \text{-N ha}^{-1} \text{ year}^{-1}$
383 or ca. 50 % of applied fertilizers in 2013. This year combined high NO_3^- concentration,
384 as previously discussed, with a humid year in which the base flow in the outlet was
385 significantly higher than in most of the remaining years (Table 3). On the other hand,
386 despite the high fertilization rates and NO_3^- concentration, the dry conditions in 2012
387 propitiated low flow and consequently $\text{NO}_3^- \text{-N}$ yield was $40 \text{ kg ha}^{-1} \text{ year}^{-1}$. Even this figure
388 implies 18% of applied N fertilisers that year. $\text{NO}_3^- \text{-N}$ yields in 2015 and 2016 were close
389 to the estimated average (81 and $53 \text{ kg ha}^{-1} \text{ year}^{-1}$ respectively), indicating that the low
390 NO_3^- concentrations observed in those years were not associated with significantly less
391 N being exported from the watershed.

392

393 According to fertilisation and soil N studies performed in several plots following typical
394 agricultural management (Litago Munarriz, 2011; INTIA and GAN, 2015b), a significant
395 oversupply of N fertilizers is applied to crops in Landazuria. In these studies, farmers did
396 not take into account either available mineral nitrogen in soils before the crop cycle (86
397 kg N ha^{-1} as a representative value) or mineral nitrogen provided by organic matter
398 mineralization ($125 - 400 \text{ kg N ha}^{-1}$ depending on site-specific conditions; INTIA and
399 GAN, 2015b). Therefore, fertilisation rates should be adjusted considering both
400 previously available N and expected organic matter mineralization rates in order to
401 minimize N leaching.

402

403 Although other forms of N may contribute to N loads in Landazuria, NO_3^- constitutes the
404 predominant form of nitrogen in eutrophic or hypertrophic waters, contributing with up to
405 90% of total N (Durand et al., 2011). In this sense, the N load estimated using NO_3^- may

406 be considered as a conservative but accurate estimation of total-N loads in this case
407 study.

408

409

410

411 *3.3. Phosphate concentration and phosphate-phosphorus yields*

412 Measured PO_4^{3-} concentrations in Landazuria outlet were below the detection limit (0.05
413 mg L^{-1}) in 63 % of collected samples (Fig. 10). In fact, PO_4^{3-} was consistently detected
414 only in specific periods, such as late summer and beginning of autumn, mainly in humid
415 years. Significant differences were observed between years (Fig. 10). The year 2015
416 presented the highest concentrations, while the years 2007, 2009 and 2010 presented
417 the lowest. It is hard to explain these pattern since no specific information about P
418 fertilization was available for this study. In addition, P dynamics may be rather complex
419 due to particulate or soluble transport pathways, and a wide range of storage and release
420 processes occurring in soils, hillslopes, groundwater or wetlands (Sharpley et al., 2013).

421

422 Monthly phosphate-phosphorus ($\text{PO}_4^{3-}\text{-P}$) loads varied widely intra- and inter-years, and
423 in general they were heavily conditioned by discharge (Table 3). For those years in which
424 a complete estimation was possible, $\text{PO}_4^{3-}\text{-P}$ yield ranged from $8 \text{ g P ha}^{-1} \text{ year}^{-1}$ in 2009,
425 to $97 \text{ g P ha}^{-1} \text{ year}^{-1}$ in the humid year 2015. It averaged $39 \pm 32 \text{ g P ha}^{-1} \text{ year}^{-1}$.

426

427

428 An important issue to consider is that, in contrast with the case of NO_3^- -N fractions of
429 total N, $\text{PO}_4^{3-}\text{-P}$ loads represent a partial estimate of total phosphorus load, since
430 particulate-P can suppose 45 – 90 % of P load in agricultural land (Merrington et al.,
431 2002). However, PO_4^{3-} (also known as soluble reactive phosphorus, SRP) is the most
432 readily bioavailable form of P, and, as a consequence, suppose the critical P pool in soils
433 and waters (Merrington et al., 2002).

434

435 Although no specific information about P fertilization was available in this study, a build-
436 up of soil P is reported to have occurred (and goes on) in agricultural soils all across
437 Europe as a consequence of over-fertilization (Daniel et al., 1998). For instance, Skhiri
438 and Dechmi (2012) reported an excess in P fertilization of about 16% of crop needs (44
439 kg P ha⁻¹ year⁻¹) in a pressurized irrigated area in the Middle Ebro Valley. However, the
440 loss of total P in the watershed outlet averaged 200 g P ha⁻¹ year⁻¹. The excessive P is
441 likely being immobilized in soils, hillslopes, streams, etc., and can imply a long-term
442 source of P in the future, or “P legacy” (Sharpley et al., 2013). This fact complicates the
443 assessment of the effects of measures to reduce P leaching from agricultural soils.

444

445 *3.4. Sediment concentration and yields*

446 A significant proportion of the collected samples (14%) presented sediment
447 concentrations below the quantification limit (0.005 g L⁻¹), coinciding with low flow
448 conditions. In general, sediment concentration in the outlet was related to discharge. The
449 higher the discharge, the higher the sediment concentration. There were some periods
450 in the year in which this pattern was not followed (Fig. 12). For instance, high sediment
451 concentrations were measured at the beginning of the irrigation season (March to June)
452 and can be related with ploughing activities before seeding of a summer crop (INTIA,
453 2017). The highest concentrations were measured when these activities coincided with
454 important precipitation events (such as in the years 2007, 2012 and 2015; Fig. 12). In
455 fact, sediment concentrations were significantly higher in the years 2007 and 2015,
456 humid years in which important precipitations occurred during the period when most of
457 the plots had been recently ploughed. The median concentration of sediment for the
458 whole study period was of 31 mg L⁻¹.

459

460 Sediment yields were greatly conditioned by extraordinary precipitation events and
461 periods (Table 3), such as that on October 4th 2013 (55 mm in a day), October 20th 2012

462 (48 mm) or September 2014, with moderate precipitations occurring after an
463 extraordinary wet summer (148 mm between June and August, twice the average in this
464 period). For those years in which a complete estimation was possible, sediment yield
465 averaged 360 kg ha⁻¹ year⁻¹, and it ranged from 42 to 1154 kg ha⁻¹ year⁻¹, being extremely
466 variable. For instance, only a few days in October 2013 accounted for 44% of the
467 sediment load for the whole study period. The importance of specific events in sediment
468 exports is a well-known fact (e.g., O'Brien et al., 2016). This is one of the main reasons
469 why long-term studies are required to adequately assess soil erosion rates (García-Ruiz
470 et al., 2015).

471

472 It is important to state that sediment concentration and yield data in Landazuria has to
473 be considered with caveats, since small wetlands are present upstream from the
474 hydrological station, adding a layer of complexity to the interpretation of sediment data.
475 The presence of these wetlands does not follow any human intervention, since they
476 developed in areas receiving irrigation return flows enriched in nutrients after the
477 transformation to irrigated land of the study area (farmer's personal communication).
478 Wetlands are recognized as effective sediment tramps (USEPA, 2015). However, they
479 may become a net source of sediment under high-flow conditions. As a comparison,
480 Aryal and Reba (2017) reported sediment concentration between 293 and 434 mg L⁻¹ for
481 an irrigated season in two watersheds in Arkansas, an order of magnitude higher than
482 that observed in Landazuria.

483

484

485 *3.5. Total dissolved solids concentrations and yields*

486 TDS concentrations in Landazuria presented a marked intra annual pattern (Fig. 14),
487 with higher values each year between late winter and early spring (the end of the non-
488 irrigated season) and lower values in the summer months (July – September, the
489 irrigated season). In addition, significant decreasing trends ($p < 0.05$) were detected for

490 every month applying the Seasonal Kendal Test (Helsel and Hirsch, 2002). Trend slopes
491 varied between -130 and -77 mg L⁻¹ year⁻¹ for the different months, with an overall value
492 of -106 mg L⁻¹ year⁻¹. In fact, the years 2015 and 2016 presented the lower TDS
493 concentrations, being significantly lower than those recorded in any other year ($p < 0.001$;
494 Fig. 14). Both the intra- and inter-annual observed pattern can be explained by the
495 influence of irrigation water in TDS concentration. The intra annual pattern is related to
496 the dilution originated by irrigation water in spring-summer months (Causapé et al.,
497 2012). On the other hand, during the study period Landazuria was a relatively recently
498 transformed irrigated area (irrigation began in the year 1999) implemented over
499 geological materials rich in gypsum and other soluble salts. Thus, the implementation of
500 irrigation mobilized salts stored in soils prior to irrigation due to prevailing semi-arid
501 climatic conditions. In addition, salts present in the geological material at the bottom of
502 Landazuria soils (marls and clays with gypsum) would be also mobilized. Consequently,
503 the amount of salts readily available for dissolution by irrigation return flows have
504 decreased with time. Similar processes have been documented in new irrigated soils
505 (Wang et al, 2012) or watersheds (Merchán et al., 2013).

506

507 Monthly TDS loads varied widely intra and inter years, and in general they were heavily
508 conditioned by discharge (Table 3). In fact, TDS loads followed a temporal distribution
509 almost identical to that of discharge. For those years in which a complete estimation was
510 possible, TDS yield ranged from 1.0 to 2.9 Mg ha⁻¹ year⁻¹ for the dry year 2012 and the
511 humid 2013, respectively. The average value was 1.8 ± 0.7 Mg ha⁻¹ year⁻¹.

512

513 An estimation of the TDS mass provided by precipitation and irrigation water was
514 obtained using its volumes and salinities. Although precipitation and irrigation water
515 presented some variability in their TDS concentration, the highest values (80 mg L⁻¹ and
516 300 mg L⁻¹, respectively; CHE, 1996) were assigned to the volumes of both components
517 of the water balance (Table 2), what implies a worst case scenario for a salt balance.

518 Under this assumption, an average of 0.35 and 1.01 Mg ha⁻¹ year⁻¹ of TDS were
519 introduced in Landazuria by precipitation and irrigation waters, respectively. Neglecting
520 minor components of the salt balance (such as salts incorporated by the crops or added
521 in fertilization, Villalobos et al., 2016), an out/in ratio of around 1.3 was obtained. Thus,
522 salts in the watershed were being washed and the feasibility of a build-up of soil salinity
523 was negligible under current irrigation and weather conditions (Thayalakumaran et al.,
524 2007).

525

526

527 *3.6. Interaction between studied variables in Landazuria*

528 Surprisingly, no significant correlation was detected between weather or agronomic
529 variables and hydrological variables (not shown). Despite the fact that Landazuria is a
530 first order watershed, a time gap does exist between any input in agricultural soils
531 (precipitation or irrigation water, fertilization...) and outputs at the watershed outlet.
532 These dynamics were not captured with the weekly resolution of the correlation analysis
533 performed in this study, and deserve further analysis using more complex time series
534 analysis. Such detailed analysis is out of the scope of the present study.

535

536 Regarding the hydrological variables themselves, the degree of correlation between
537 discharge and concentrations was, in general, low or non-significant (Table 4). This may
538 be a consequence of the amount of different processes involved and the different initial
539 conditions for rainfall-runoff events, fertilisation season, etc. Only subtle significant
540 relationships were detected between discharge and concentration of NO₃⁻ (-0.18) and
541 sediment (+0.19). However, stronger correlations were detected between discharge and
542 loads: high correlation coefficients were obtained for TDS and NO₃⁻-N loads, while the
543 correlation coefficients between PO₄³⁻-P and sediment loads were lower. Isidoro et al.
544 (2003) also detected low correlation between discharge and water quality parameters in
545 other irrigated area of the Middle Ebro Valley.

546

547 In general, low correlation coefficients were obtained between the different water quality
548 variables (Table 4), indicating different processes controlling each of them. Edwards and
549 Withers (1998) also reported the differences in hydrological processes controlling each
550 pollutant dynamics in agricultural watersheds in the UK. In Landazuria, high
551 concentration of one parameter did not coincide with high concentrations in any other
552 one. Some degree of correlation was detected between the soluble constituents (NO_3^-
553 vs. TDS = +0.41) and were probably due to dilution effects after strong precipitation
554 events or under increased base flow conditions in the irrigated season. In contrast, higher
555 correlation coefficients were obtained between the loads of different constituents. NO_3^- -
556 N and TDS loads were strongly related between them (+0.93) and more weakly related
557 to the remaining constituents (+0.34 - +0.47). The relation between PO_4^{3-} -P and
558 sediment loads was lower (+0.53). Although some studies have estimated that up to 90
559 % of P load occurs in particulate forms (Merrington et al., 2002), high P concentration in
560 fine sediment mobilized under high flow conditions makes PO_4^{3-} to desorb from the
561 sediment into the water (Gibson, 1997). As a consequence, PO_4^{3-} -P load (and PO_4^{3-} -
562 concentrations) tended to increase with sediment loads.

563

564 For soluble constituents, small correlation coefficients were obtained between each
565 water quality variable (i.e., concentration) and the constituent load (Table 4), indicating
566 that the higher loads did not coincide with the higher concentrations (+0.18 for NO_3^- vs.
567 NO_3^- -N Load, and +0.22 for TDS vs. TDS Load). In contrast, a strong correlation was
568 founded between concentration and loads (+0.83 and +0.91 for P and sediments,
569 respectively) indicating that high concentrations coincided with high loads events.

570

571 The observed relationships support the fact that different processes control the
572 behaviour of these pollutants: NO_3^- and TDS represent highly soluble components that
573 were easily mobilized by flowing water, either under base flow or high flow conditions.

574 On the other hand, PO_4^{3-} and sediment have low solubility and therefore their
575 concentration was linked to specific high flow events that eroded soil particles in the
576 fields, banks or bare soil areas (Aryal and Reba, 2017).

577

578 *3.7. Agricultural pollution dynamics in irrigated areas in contrast with rainfed areas*

579 Irrigated agriculture is mostly located in semi-arid areas, where the climatic conditions
580 do not allow to achieve competitive agronomic production. Regions with Mediterranean
581 climate, where the higher temperatures of summer coincide with the dry season,
582 constitute a typical example (Ryan et al., 2009). Soils in semi-arid regions typically have
583 a particular set of characteristics, such as lower organic matter content or higher pH and
584 carbonate contents, in relation with soils in more humid regions (Brady and Weil, 2008;
585 Ryan et al., 2009). In addition, climatic conditions imply, in general, low yield in these
586 regions, especially for summer crops. As a consequence, winter crops such as wheat or
587 barley are predominant (Ryan et al., 2009), and its productivity rely on climatic
588 conditions. All these characteristics are present in Landazuria as reported in section 2.1:
589 semi-arid conditions, especially with hot and dry summers; predominance of alkaline and
590 calcareous soils; fallow-winter cereal as the typical rotation in the rainfed area, with low
591 yield (ca. 2.5 and 4 times less yield in the rainfed surface in comparison with the irrigated
592 one for barley and wheat, respectively).

593

594 Thus, irrigation in semi-arid soils produce a significant modification in the moisture
595 conditions and the soil water balance, especially during summer when most of irrigation
596 is applied. As a consequence, several properties of the soil are modified. For instance,
597 the cultivation of more productive crops implies an increase in crop residues, and
598 therefore organic matter in these soils (Apesteguía et al., 2015). In addition, summer
599 high temperatures in combination with moisture provided by irrigation significantly
600 increase organic matter mineralization rates (Arroita et al., 2013).

601

602 The water balance in watersheds with a significant proportion of irrigated surface is
603 severely modified. However, the casuistic in watersheds receiving irrigation water from
604 external sources is quite different to those in which irrigation water is obtained through
605 dams, diversions or aquifers from the watershed itself. Indeed, in watersheds with
606 external sources of irrigation water, excessive irrigation applied over the crop needs or
607 summer rainfalls in already wet soils facilitates water leaching and, thus, increases
608 aquifer recharge and/or baseflow contributions to nearby streams (e.g., Cameira et al.,
609 2003; Sánchez-Pérez et al., 2003). In contrast, decreased flow of rivers or depleted
610 aquifers are the main consequences when the irrigation water is obtained within the
611 analysed watershed (e.g., Scanlon et al., 2007). In the rest of this discussion, we focus
612 in the environmental issues in watersheds receiving irrigation water from external
613 sources, such as Landazuria.

614

615 In fact, Landazuria exemplifies this modified water balance, with stable discharge
616 throughout the year. In contrast, rainfed watersheds in Navarre, even in more humid
617 areas, presented summer discharge values considerably lower than those observed in
618 Landazuria (Casalí et al., 2008, 2010). Other studies have reported the severe
619 modification of the water balance in irrigated areas (Barros et al., 2011; Andrés and
620 Cuchí, 2014). In fact, in a watershed with similar conditions to that observed in
621 Landazuria, Merchán et al. (2013) reported the shift from an ephemeral stream to a
622 permanent one after the implementation of irrigation.

623

624 Water balance modification in semi-arid soils and watersheds has a huge relevance on
625 its environmental impact since it controls pollutants exports to water bodies. Nitrate, as
626 a readily soluble constituent of soil water, is particularly affected by modifications in the
627 water balance. In fact, despite similar fertilization rates, nitrate-N yield from Landazuria
628 irrigated surface (ca. 70 kg ha⁻¹ year⁻¹) was higher than that in rainfed winter cereal
629 watersheds in Navarre (16-37 kg ha⁻¹ year⁻¹, Casalí et al., 2008). While nitrate-N yield

630 in rainfed cereal watersheds occurred mostly during winters and was negligible during
631 summer, in Landazuria it occurred throughout the year, as a consequence of the
632 increased summer discharge due to irrigation return flows. In addition, more productive
633 crops are usually grown under higher water availability (e.g., Gaydon et al., 2012), raising
634 significantly the expected production and, consequently, fertilization rates. In Landazuria,
635 the irrigated surface presented: (1) maize and vegetables such as tomato or onion as
636 dominant crops; (2) a significant proportion of fields with two crops per year; and (3) an
637 average fertilization rate seven times higher than that in the rainfed surface, where winter
638 cereals were the dominant crops and cultivation took place one out of two years (Table
639 1).

640

641 A simple balance between average fertilization rates and nitrate-N yield in Landazuria
642 suggested that around 35 % of N fertilizer being lost in drainage. Although a great range
643 of variation is reported in the literature, similar figures have been reported in irrigated
644 areas in Spain or Australia (Barros et al., 2012b; Thorburn et al., 2011). The variation in
645 N losses fraction in drainage have been related to differences in terms of soils, crops
646 grown and agricultural management. As extreme values, no significant N loss through
647 drainage have been reported in irrigated clayey soils (Arauzo and Valladolid, 2013)
648 whereas losses of up to 77 % of applied N fertilizer have been reported for humid areas
649 with well-drained sandy soils intensively cropped in central Wisconsin, US (Kraft and
650 Stites, 2003).

651

652 Irrigation by itself does not imply higher nitrate leaching. In fact, in more humid climates
653 nitrate leaching is extremely variable, depending on climatic, soils or management
654 conditions. For instance, in rainfed soils over terraces in Nepal, nitrate-N leaching ranged
655 between 0-64 kg ha⁻¹ year⁻¹. In general, it was observed that the more fertilizer received,
656 the more leaching (Pilbeam et al., 2004). In long-term monitored watersheds, the
657 agricultural land exported 6-32 and 10-40 kg N ha⁻¹ year⁻¹ in Sweden and Estonia,

658 respectively (Kyllmar et al., 2014; Iñal et al., 2014). However, even under humid
659 conditions, nitrate-N leaching is expected to increase in irrigated and heavily fertilized
660 areas (Kraft and Stites, 2003).

661

662 Due to the aforementioned calcareous soils, phosphate-P leaching is not particularly
663 enhanced by irrigation in semi-arid regions, as the P-fixing capacity of these soils is
664 considerable. For instance, phosphate-P yield in Landazuria (39 g ha⁻¹ year⁻¹) was in
665 the order of magnitude of that estimated for rainfed winter cereal watersheds in Navarre
666 (52 g ha⁻¹ year⁻¹; Casalí et al., 2008). However, a watershed with more humid
667 conditions, neutral soils and manure application presented higher phosphate-P yield
668 (120-250 g ha⁻¹ year⁻¹; Casalí et al., 2010). According to Withers and Sharpley (1995),
669 soluble P makes up a greater component of P in runoff from sites that frequently receive
670 applications of organic manures and slurries. In Alberta, Olson et al. (2010) related risk
671 of P leaching with application rates of manure in irrigated soils. Thus, factors such as soil
672 pH or organic fertilization, rather than irrigation, controls phosphate-P leaching in
673 agricultural watersheds. Total P, in contrast, is greatly related to erosion and sediment
674 losses from agricultural fields (Merrington et al., 2002). In fact, a significant relationship
675 was detected between phosphate-P and sediment loads in Landazuria (section 3.6,
676 Table 4).

677

678 Irrigation water quality and temperature, application rates or irrigation systems make
679 irrigation-induced erosion a phenomenon quite different from rainfed erosion (Sojka et
680 al., 2008). Unfortunately, there is scarce data about irrigation-induced erosion with a wide
681 range of reported values in the available studies. Indeed, both rainfall- and irrigation-
682 induced erosion contribute to soil loss in irrigated areas. Despite this, sediment yield in
683 Landazuria (360 kg ha⁻¹ year⁻¹) was in the lower range of those estimated for the rainfed
684 winter cereal watersheds in Navarre (230-1350 kg ha⁻¹ year⁻¹). The presence of
685 wetlands (section 3.4) and the lower slopes in Landazuria (3–5%) in comparison with the

686 rainfed winter cereal watersheds (7–30%) were the most likely explanations for the
687 different behaviour. Although no much data was available for comparison, in a terraced
688 surface irrigated area of the Middle Ebro Valley, the sediment yield estimation was similar
689 to that obtained in this study (ca. 300 kg ha⁻¹ year⁻¹; García-Ruiz, 2010).

690

691 The clearing of natural vegetation for rainfed crop production, especially in arid to semi-
692 arid areas, is reported to increase mobilization of salts stored in the vadose zone over
693 millennia due to an increases in soil water leaching (Scanlon et al., 2007). The shift to
694 irrigated agriculture normally enhances this process due to an increase in available water
695 for leaching (Merchán et al., 2015a). The rate at which salts are washed out depends
696 mainly on the available amount of salts and the water leached. The amount of available
697 salts depends on the geological materials and the “recent” (in geological terms) history
698 of the soil. For instance, in many places of Australia, a great degree of irrigation induced
699 leaching of salts is reported as a consequence of prevailing climatic conditions during
700 recent times (Duncan et al., 2008). On the other hand, the amount of irrigation return
701 flows is controlled by the lack or existence of irrigation, and the system and/or
702 management of irrigation carried out. In this sense, higher irrigation efficiencies have
703 been related to lower salt loading (e.g., García-Garizábal and Causapé, 2010; Merchán
704 et al., 2015c).

705

706 3.8. Overview of management practices to mitigate the environmental impact of irrigated
707 areas in receiving water bodies

708 In a meta-analysis conducted over irrigated soils studies (Quemada et al., 2013),
709 improved water and fertilization management were reported as the most effective
710 practices in reducing nitrate leaching, with average decreases of 58 % and 39 %,
711 respectively. Alternative practices such as use of cover crops or improved fertilization
712 technologies presented, in general, lower reductions in leaching (Quemada et al., 2013).
713 This behaviour is also detected at the watershed scale. For instance, improved surface

714 irrigation management decreased the yield of soluble constituents such as nitrate and
715 salts (Barros et al. 2012a,b; García-Garizábal et al., 2012, 2014). In general, sprinkler
716 irrigation presented lower yields of nitrate and salts than surface irrigation (Merchán et
717 al., 2015c).

718

719 Nitrogen fertilization rates should be adapted to the necessities of the crops, both in total
720 amount applied and temporal distribution (Quemada et al., 2013), as these measures
721 significantly reduced nitrate leaching with minimal impact on yield. Reduced fertilization
722 rates decreased leaching but also yield, while fertigation did not always reduce leaching
723 and it decreased in ca. 6 % yield (Quemada et al., 2013). In addition, the mineral N
724 available at the beginning of the crop cycle along with that provided by organic matter
725 mineralisation should be accounted for (Cameira et al., 2003; INTIA, 2017). Regarding
726 phosphorus, applications in manure and slurries, rather than irrigation, usually controls
727 phosphate leaching (e.g., Olson et al., 2010).

728

729 Since irrigated areas are subject to both rainfall- and irrigation-induced erosion, a wide
730 range of management practices can be applied to decrease soil loss and sediment yield
731 in irrigated areas. Apart from those typical of rainfed areas, practices such as improved
732 water management, tailwater elimination/reuse or low-pressure wide-area sprays are
733 recommended to curb irrigation-induced erosion (Sojka et al., 2008). For instance,
734 Campo-Bescós et al. (2015) reported the beneficial effects of improved irrigation
735 management and vegetative filter strips in the sediment exported in furrow-irrigated
736 fields.

737

738 Although detailed information about irrigation management was not available for
739 Landazuria, the higher discharge in summer with respect to winter suggest there is some
740 margin of improvement in irrigation management that would probably decrease the
741 environmental impact in terms of nitrate and salt yield. N fertilisation management clearly

742 needs to be optimized since the average loss of N in the watershed was about one third
743 of N applied in fertilisers. Phosphate and sediment yield were not considered especially
744 problematic in this study, probably due to the low use of organic fertilization and the
745 presence of semi-natural wetlands and relatively low slopes in Landazuria.

746

747

748

749 **4. Conclusion**

750 In this study, a detailed and long-term dataset collected in the watershed outlet is used
751 to estimate pollutant loads and behaviour in an area under pressurized irrigation typical
752 in north-eastern Spain. While the inclusion of several variables provides an integrated
753 picture of the studied system, the length of the obtained record (including both normal
754 and extreme climatic and agronomic conditions) provided reliable estimates on the
755 environmental effects of irrigation in this particular study case. This estimates are then
756 contextualized in the broad field of agricultural pollution of water bodies.

757

758 Irrigation modified the water regime of Landazuria stream, providing a higher base flow
759 during summer months along with a relatively constant discharge throughout the year.
760 Median concentration in daily water samples collected at the watershed outlet for ten
761 agricultural years were: NO_3^- , 185 mg L⁻¹; PO_4^{3-} (or SRP) <0.05 mg L⁻¹; sediment, 31 mg
762 L⁻¹; TDS, 2284 mg L⁻¹. Estimated annual yields averaged 39 kg NO_3^- -N ha⁻¹, 39 g PO_4^{3-} -
763 P ha⁻¹, 360 kg sediment ha⁻¹ and 1.8 Mg TDS ha⁻¹. For both concentration and loads, a
764 high degree of intra- and inter-annual variation was observed, conditioned by different
765 weather conditions, the behaviour of highly soluble constituents (NO_3^- and TDS) or
766 particulate driven constituents (sediment and PO_4^{3-}), and agronomic management. In
767 these sense, NO_3^- and TDS behaved conservatively with respect to water whereas PO_4^{3-}
768 and sediment dynamics were related to high flow events.

769

770 Our results were put into a broader context, especially regarding differences between
771 rainfed and irrigated agricultural systems in semi-arid areas receiving irrigation water
772 from external sources (neighbouring watersheds). In this sense, irrigation water
773 supposes an additional water input, increasing aquifer recharge and baseflow of nearby
774 streams. It implies a shift to more productive and N-fertilizer demanding crops, and
775 consequently tends to increase nitrate leaching in these soils. In addition, salt stored in
776 the vadose zone are mobilized by irrigation return flows, increasing salinity in
777 downstream water bodies. Phosphorus fixing capacity of semi-arid soils implies that
778 phosphate leaching is not particularly enhanced by irrigation, while erosion and sediment
779 yield from irrigated areas depends greatly in site-specific characteristics.

780

781

782 Given the high variability in both concentration and loads imposed by the precipitation
783 regime and agronomic management, long-term monitoring in well characterized
784 agricultural watersheds is essential to understand the dynamics of pollution processes.
785 In addition, it can provide an accurate estimation of the pollution “base level”, which is
786 needed in order to assess the effects of preventive or corrective measures undertaken
787 to improve the water quality downstream.

788

789

790 Acknowledgements

791 This work was possible thanks to the monitoring network of the Government of Navarre and the
792 detailed agronomic information collected by INTIA in the framework of the LIFE-Nitrates Project
793 (LIFE+10 ENV/ES/478; www.life-nitratos.eu). In addition, it benefited from the Research Project
794 CGL2015-64284-C2-1-R (Ministerio de Economía y Competitividad). Daniel Merchán
795 acknowledges the support of the “Juan de la Cierva - Formación” grant with code FJCI-2015-
796 24920. We appreciate the collaboration and information provided by the farmers and the Irrigation
797 Authority (Comunidad de Regantes El Ferial) in Landazuria.

798

799 **References**

- 800 Allen, R.G., Pereira, L.S., Raes, D., Smith, M., 1998. Crop Evapo-Transpiration: Guidelines for
801 Computing Crop Water Requirements. United Nations Food and Agriculture
802 Organization, Rome, Italy (FAO Irrigation and Drainage Paper No. 56).
- 803 Andrés, R., Cuchí, J.A., 2014. Analysis of sprinkler irrigation management in the LASESA
804 district, Monegros (Spain). *Agric. Water Manag.* 131, 95–107.
805 doi:10.1016/j.agwat.2013.09.016
- 806 Apesteguía, M., Virto, I., Orcaray, L., Enrique, A., Bescansa, P., 2015. Effect of the Conversion
807 to Irrigation of Semiarid Mediterranean Dryland Agroecosystems on Soil Carbon
808 Dynamics and Soil Aggregation. *Arid Land Res. Manag.* 0, 1-16.
809 doi:10.1080/15324982.2015.1016245
- 810 Arauzo, M., Valladolid, M., 2013. Drainage and N-leaching in alluvial soils under agricultural
811 land uses: Implications for the implementation of the EU Nitrates Directive. *Agric.
812 Ecosyst. Environ.* 179, 94–107. doi:10.1016/j.agee.2013.07.013
- 813 Arroita, M., Causapé, J., Comín, F.A., Díez, J., Jimenez, J.J., Lacarta, J., Lorente, C., Merchán,
814 D., Muñiz, S., Navarro, E., Val, J., Elozegi, A., 2013. Irrigation agriculture affects
815 organic matter decomposition in semi-arid terrestrial and aquatic ecosystems. *J.
816 Hazard. Mater.* 263, 139–145. doi:10.1016/j.jhazmat.2013.06.049
- 817 Aryal, N., Reba, M.L., 2017. Transport and transformation of nutrients and sediment in two
818 agricultural watersheds in Northeast Arkansas. *Agric. Ecosyst. Environ.* 236, 30–42.
819 doi:10.1016/j.agee.2016.11.006
- 820 Barros, R., Isidoro, D., Aragüés, R., 2011. Long-term water balances in La Violada Irrigation
821 District (Spain): II. Analysis of irrigation performance. *Agric. Water Manag.* 98, 1569–
822 1576. doi:10.1016/j.agwat.2011.04.014
- 823 Barros, R., Isidoro, D., Aragüés, R., 2012a. Three study decades on irrigation performance and
824 salt concentrations and loads in the irrigation return flows of La Violada irrigation district
825 (Spain). *Agric. Ecosyst. Environ.* 151, 44–52. doi:10.1016/j.agee.2012.02.003
- 826 Barros, R., Isidoro, D., Aragüés, R., 2012b. Irrigation management, nitrogen fertilization and
827 nitrogen losses in the return flows of La Violada irrigation district (Spain). *Agric.
828 Ecosyst. Environ.* 155, 161–171. doi:10.1016/j.agee.2012.04.004
- 829 Brady, N.C., Weil, R.R., 2008. *The Nature and Properties of Soils*, 14th ed. Pearson Education,
830 Inc., Upper Saddle River, New Jersey, USA.
- 831 Burt, T.P., Howden, N.J.K., Worrall, F., Whelan, M.J., 2010. Long-term monitoring of river water
832 nitrate: how much data do we need? *J. Environ. Monit.* 12, 71–79.
833 doi:10.1039/B913003A
- 834 Cameira, M.R., Fernando, R.M., Pereira, L.S., 2003. Monitoring water and NO₃-N in irrigated
835 maize fields in the Sorraia Watershed, Portugal. *Agric. Water Manag.* 60, 199–216.
836 doi:10.1016/S0378-3774(02)00175-0

837 Campo-Bescós, M.A., Muñoz-Carpena, R., Kiker, G.A., Bodah, B.W., Ullman, J.L., 2015.
838 Watering or buffering? Runoff and sediment pollution control from furrow irrigated fields
839 in arid environments. *Agric. Ecosyst. Environ.* 205, 90–101.
840 doi:10.1016/j.agee.2015.03.010

841 Casalí, J., Gastesi, R., Álvarez-Mozos, J., De Santisteban, L.M., Del Valle de Lersundi, J.,
842 Giménez, R., Larrañaga, A., Goñi, M., Agirre, U., Campo, M.A., López, J.J., Donézar,
843 M., 2008. Runoff, erosion, and water quality of agricultural watersheds in central
844 Navarre (Spain). *Agric. Water Manag.* 95, 1111–1128. doi:10.1016/j.agwat.2008.06.013

845 Casalí, J., Giménez, R., Díez, J., Álvarez-Mozos, J., Del Valle de Lersundi, J., Goñi, M., Campo,
846 M.A., Chahor, Y., Gastesi, R., López, J., 2010. Sediment production and water quality
847 of watersheds with contrasting land use in Navarre (Spain). *Agric. Water Manag.* 97,
848 1683–1694. doi:10.1016/j.agwat.2010.05.024

849 Causapé, J., Merchán, D., Abrahão, R., García-Garizábal, I., 2012. Hydrological changes in
850 Lerma creek (Zaragoza) after the implementation of irrigation. *Cuadernos de*
851 *Investigación Geográfica* 38(2), 91-106. doi: 10.18172/cig.1284

852 CHE (Confederación Hidrográfica del Ebro – Ministerio de Medio Ambiente), 1996. Diagnóstico
853 y gestión ambiental de embalses en el ámbito de la cuenca hidrográfica del Ebro:
854 Embalse de Yesa. *Limnos*, 19pp.

855 CHE, 2017. “Datos básicos de la Confederación Hidrográfica del Ebro”. Available at:
856 www.chebro.es/contenido.visualizar.do?idContenido=37945&idMenu=2167 [Accessed
857 May 2017, in Spanish]

858 CR-El Ferial, 2017. “Históricos de Cultivos en la Comunidad de Regantes El Ferial”. Available
859 at: <http://www.riegosdenavarra.com/agroind2/bard4.htm> [Accessed March 2017, in
860 Spanish]

861 Daniel, T.C., Sharpley, A.N., Lemunyon, J.L., 1998. Agricultural Phosphorus and
862 Eutrophication: A Symposium Overview. *J. Environ. Qual.* 27, 251–257.
863 doi:10.2134/jeq1998.00472425002700020002x

864 DDRMAAL (Departamento de Desarrollo Rural, Medio Ambiente y Administración Local –
865 Gobierno de Navarra), 2017. “Estadísticas agrícolas”. *Negociado de Estadística*.
866 Available at:
867 [http://www.navarra.es/home_es/Temas/Ambito+rural/Vida+rural/Observatorio+agrario/
868 Agricola/Informacion+estadistica/](http://www.navarra.es/home_es/Temas/Ambito+rural/Vida+rural/Observatorio+agrario/Agricola/Informacion+estadistica/) [Accessed March 2017, Spanish]

869 DOPTC (Departamento de Obras Públicas, Transportes y Comunicaciones – Gobierno de
870 Navarra), 2003a. “Cartografía Geológica de Navarra, Escala 1:25.000, 244-II (Rada)”.
871 Geological map and report. 119 pp. Available at:
872 http://www.navarra.es/home_es/Temas/Territorio [Accessed March 2017, Spanish]

873 DOPTC, 2003b. “Cartografía Geológica de Navarra, Escala 1:25.000, 244-IV (Arguedas)”.
874 Geological map and reports. 120 pp. Available at:
875 http://www.navarra.es/home_es/Temas/Territorio [Accessed March 2017, Spanish]

876 Duncan, R.A., Bethune, M.G., Thayalakumaran, T., Christen, E.W., McMahon, T.A., 2008.
877 Management of salt mobilisation in the irrigated landscape - A review of selected
878 irrigation regions. *J. Hydrol.* 351, 238–252. doi:10.1016/j.jhydrol.2007.12.002

879 Durand, P., et al., 2011. Nitrogen processes in aquatic ecosystems, in: Sutton, M.A., Howard,
880 C.M., Erisman, J.W., Billen, G., Bleeker, A., Grennfelt, P., van Grinsven, H., Grizzetti,
881 B. (Eds.), *The European Nitrogen Assessment: Sources, Effects and Policy*
882 *Perspectives*. Cambridge University Press, Cambridge, UK, pp. 126–146.

883 Edwards, A.C., Withers, P.J.A., 2008. Transport and delivery of suspended solids, nitrogen and
884 phosphorus from various sources to freshwaters in the UK. *J. Hydrol.* 350, 144–153.
885 doi:10.1016/j.jhydrol.2007.10.053

886 Elmi, A.A., Madramootoo, C., Egeh, M., Hamel, C., 2004. Water and fertilizer nitrogen
887 management to minimize nitrate pollution from a cropped soil in southwestern Quebec,
888 Canada. *Water. Air. Soil Pollut.* 151, 117–134.
889 doi:10.1023/B:WATE.0000009910.25539.75

890 FAO, 2003a. *Unlocking the Water Potential of Agriculture*. United Nations Food and Agriculture
891 Organization, Rome, Italy.

892 FAO, 2003b. *World Agriculture Towards 2015/2030. A FAO Perspective*. United Nations Food
893 and Agriculture Organization, Rome, Italy.

894 FAO, 2013. *FAO Statistical Year Book 2013, World Food and Agriculture*. United Nations Food
895 and Agriculture Organization, Rome, Italy.

896 García-Garizábal, I., Causapé, J., 2010. Influence of irrigation water management on the
897 quantity and quality of irrigation return flows. *J. Hydrol.* 385, 36–43.
898 doi:10.1016/j.jhydrol.2010.02.002

899 García-Garizábal, I., Causapé, J., Abrahão, R., 2012. Nitrate contamination and its relationship
900 with flood irrigation management. *J. Hydrol.* 442–443, 15–22.
901 doi:10.1016/j.jhydrol.2012.03.017

902 García-Ruiz, J.M., 2010. The effects of land uses on soil erosion in Spain: A review. *Catena* 81,
903 1–11. doi:10.1016/j.catena.2010.01.001

904 García-Ruiz, J.M., Beguería, S., Nadal-Romero, E., González-Hidalgo, J.C., Lana-Renault, N.,
905 Sanjuán, Y., 2015. A meta-analysis of soil erosion rates across the world.
906 *Geomorphology* 239, 160–173. doi:10.1016/j.geomorph.2015.03.008

907 Gaydon, D.S., Meinke, H., Rodriguez, D., 2012. The best farm-level irrigation strategy changes
908 seasonally with fluctuating water availability. *Agric. Water Manag.* 103, 33–42.
909 doi:10.1016/j.agwat.2011.10.015

910 Gybson, C.E., 1997. The dynamics of phosphorus in freshwater and marine environments, in:
911 Tunney, H., Carotn, O.T., Brookes, P.C., Johnston, A.E. (Eds.), *Phosphorus Losses*
912 *from Soil to Water*. CAB International, Wallingford, UK, pp. 119–135.

913 Government of Navarre, 2005. “Memoria del Mapa de Suelos de la Cuenca Agraria
914 Experimental de Landazuria, Escala 1/25.000”. Soils map and report. 97 pp.
915 [Unpublished, Spanish]

916 Government of Navarre, 2017. PAC Declarations, Personal Communication.

917 Helsel, D.R., Hirsch, R.M., 2002. Statistical methods in water resources. US Geological Survey,
 918 Reston, VA. 510 pp.

919 IDENA (Infraestructura de Datos Espaciales de Navarra – Gobierno de Navarra), 2017.
 920 Available at: <https://idena.navarra.es/Portal/Descargar> [Accessed March 2017,
 921 Spanish]

922 Iital, A., Klõga, M., Pihlak, M., Pachel, K., Zahharov, A., Loigu, E., 2014. Nitrogen content and
 923 trends in agricultural catchments in Estonia. *Agric. Ecosyst. Environ.* 198, 44–53.
 924 doi:10.1016/j.agee.2014.03.010

925 INTIA (Instituto Navarro de Tecnologías e Infraestructuras Agroalimentarias), 2017. Agronomic
 926 management, farmers face-to-face inquiries, database. Proyecto LIFE Nitratos
 927 (LIFE+10 ENV/ES/478). [Unpublished, Spanish]

928 INTIA and GAN (Instituto Navarro de Tecnologías e Infraestructuras Agroalimentarias and
 929 Gestión Ambiental de Navarra), 2015a. “Buenas Prácticas Agrarias: Directrices”.
 930 Proyecto LIFE Nitratos (LIFE+10 ENV/ES/478). Ed: INTIA, GAN y Fundación CRANA –
 931 Gobierno de Navarra. 36 pp. Available at:
 932 <http://www.life-nitratos.eu/index.php/es/documentos/documentos-del-proyecto>
 933 [Accessed March 2017, Spanish]

934 INTIA and GAN, 2015b. “Informe de los Seminarios Finales”. Proyecto LIFE Nitratos (LIFE+10
 935 ENV/ES/478). Ed: INTIA, GAN y Fundación CRANA – Gobierno de Navarra. 275 pp.
 936 Available at:
 937 <http://www.life-nitratos.eu/index.php/es/documentos/documentos-del-proyecto>
 938 [Accessed March 2017, Spanish]

939 Isidoro, D., Quílez, D., Aragüés, R., 2003. Sampling strategies for the estimation of salt and
 940 nitrate loads in irrigation return flows: La Violada Gully (Spain) as a case study. *J.*
 941 *Hydrol.* 271, 39–51. doi:10.1016/S0022-1694(02)00324-4

942 Kraft, G.J., Stites, W., 2003. Nitrate impacts on groundwater from irrigated-vegetable systems in
 943 a humid north-central US sand plain. *Agric. Ecosyst. Environ.* 100, 63–74.
 944 doi:10.1016/S0167-8809(03)00172-5

945 Kyllmar, K., Stjernman Forsberg, L., Andersson, S., Mårtensson, K., 2014. Small agricultural
 946 monitoring catchments in Sweden representing environmental impact. *Agric. Ecosyst.*
 947 *Environ.* 198, 25–35. doi:10.1016/j.agee.2014.05.016

948 Letey, J., Hoffman, G.J., Hopmans, J.W., Grattan, S.R., Suarez, D., Corwin, D.L., Oster, J.D.,
 949 Wu, L., Amrhein, C., 2011. Evaluation of soil salinity leaching requirement guidelines.
 950 *Agric. Water Manag.* 98, 502–506. doi:10.1016/j.agwat.2010.08.009

951 Litago Munarriz, J., 2011. “Balances de nitrógeno y pérdidas de nitratos por lixiviación en
 952 parcelas de regadío”. BSc Ag. Eng. Dissertation, Universidad Pública de Navarra, 123
 953 pp. [Spanish]

954 MAPAMA (Ministerio de Agricultura y Pesca, Alimentación y Medio Ambiente – Gobierno de
 955 España), 2017. Gestión sostenible de regadíos.

956 <http://www.mapama.gob.es/es/desarrollo-rural/temas/gestion-sostenible-regadios/>
957 [Accessed March 2017, Spanish]

958 Merchán, D., Causapé, J., Abrahão, R., 2013. Impact of irrigation implementation on hydrology
959 and water quality in a small agricultural basin in Spain. *Hydrol. Sci. J.* 58, 1400–1413.
960 doi:10.1080/02626667.2013.829576

961 Merchán, D., Auqué, L.F., Acero, P., Gimeno, M.J., Causapé, J., 2015a. Geochemical
962 processes controlling water salinization in an irrigated basin in Spain: Identification of
963 natural and anthropogenic influence. *Sci. Total Environ.* 502, 330–343.
964 doi:10.1016/j.scitotenv.2014.09.041

965 Merchán, D., Causapé, J., Abrahão, R., García-Garizábal, I., 2015b. Assessment of a newly
966 implemented irrigated area (Lerma Basin, Spain) over a 10-year period. I: Water
967 balances and irrigation performance. *Agric. Water Manag.* 158, 277–287.
968 doi:10.1016/j.agwat.2015.04.016

969 Merchán, D., Causapé, J., Abrahão, R., García-Garizábal, I., 2015c. Assessment of a newly
970 implemented irrigated area (Lerma Basin, Spain) over a 10-year period. II: Salts and
971 nitrate exported. *Agric. Water Manag.* 158, 288–296. doi:10.1016/j.agwat.2015.04.019

972 Merrington, G., Winder, L., Parkinson, R., Redman, M., 2002. *Agricultural Pollution,*
973 *Environmental problems and practical solutions.* Spon's Environmental Science and
974 *Engineering Series.* Spon Press, London and New York.

975 Nielsen, D.L., Brock, M.A., Rees, G.N., Baldwin, D.S., 2003. Effects of increasing salinity on
976 freshwater ecosystems in Australia. *Aust. J. Bot.* 51(6), 655–665. doi:10.1071/BT02115

977 Novotny, V., 1999. Diffuse pollution from agriculture - a worldwide outlook. *Water Sci. Technol.*
978 39 (3), 1–13.

979 O'Brien, K.R., Weber, T.R., Leigh, C., Burford, M.A., 2016. Sediment and nutrient budgets are
980 inherently dynamic: evidence from a long-term study of two subtropical reservoirs.
981 *Hydrol. Earth Syst. Sci.* 20, 4881–4894. doi:10.5194/hess-20-4881-2016

982 Olson, B.M., Bremer, E., McKenzie, R.H., Bennett, R., 2010. Phosphorus accumulation and
983 leaching in two irrigated soils with incremental rates of cattle manure. *Can. J. Soil Sci.*
984 90, 355–362. doi:10.4141/CJSS09025

985 Playán, E., Salvador, R., Faci, J.M., Zapata, N., Martínez-Cob, A., Sánchez, I., 2005. Day and
986 night wind drift and evaporation losses in sprinkler solid-sets and moving laterals. *Agric.*
987 *Water Manag.* 76, 139–159. doi:10.1016/j.agwat.2005.01.015

988 Pilbeam, C.J., Gregory, P.J., Munankarmy, R.C., Tripathi, B.P., 2004. Leaching of nitrate from
989 cropped rainfed terraces in the mid-hills of Nepal. *Nutr. Cycl. Agroecosystems* 69, 221–
990 232. doi:10.1023/B:FRES.0000035194.15958.e0

991 Quemada, M., Baranski, M., Nobel-de Lange, M.N.J., Vallejo, A., Cooper, J.M., 2013. Meta-
992 analysis of strategies to control nitrate leaching in irrigated agricultural systems and
993 their effects on crop yield. *Agric. Ecosyst. Environ.* 174, 1–10.
994 doi:10.1016/j.agee.2013.04.018

995 Rivett, M.O., Buss, S.R., Morgan, P., Smith, J.W.N., Bemment, C.D., 2008. Nitrate attenuation
996 in groundwater: A review of biogeochemical controlling processes. *Water Res.* 42,
997 4215–4232. doi:10.1016/j.watres.2008.07.020

998 Ryan, J., Ibrikci, H., Sommer, R., McNeill, A., 2009. Nitrogen in rainfed and Irrigated Cropping
999 Systems in the Mediterranean Region. *Advances in Agronomy* 104, 53-106.
1000 doi:10.1016/S0065-211-(09)04002-4

1001 Sánchez Pérez, J.M., Antigüedad, I., Arrate, I., García-Linares, C., Morell, I., 2003. The
1002 influence of nitrate leaching through unsaturated soil on groundwater pollution in an
1003 agricultural area of the Basque country: A case study. *Sci. Total Environ.* 317, 173–187.
1004 doi:10.1016/S0048-9697(03)00262-6

1005 Scanlon, B.R., Jolly, I., Sophocleous, M., Zhang, L., 2007. Global impacts of conversions from
1006 natural to agricultural ecosystems on water resources: Quantity versus quality. *Water*
1007 *Resour. Res.* 43. doi:10.1029/2006WR005486

1008 Sharpley, A., Jarvie, H.P., Buda, A., May, L., Spears, B., Kleinman, P., 2014. Phosphorus
1009 Legacy: Overcoming the Effects of Past Management Practices to Mitigate Future
1010 Water Quality Impairment. *J. Environ. Qual.* 42, 1308–1326.
1011 doi:10.2134/jeq2013.03.0098

1012 Skhiri, A., Dechmi, F., 2012. Impact of sprinkler irrigation management on the Del Reguero river
1013 (Spain) II: Phosphorus mass balance. *Agric. Water Manag.* 103, 130–139.
1014 doi:10.1016/j.agwat.2011.11.004

1015 Sojka, E.E., Bjerneberg, D.L., Strelkoff, T.S., 2008. Irrigation-Induced Erosion. In: *Irrigation of*
1016 *Agricultural Crops*, 2nd ed. *Agronomy Monograph* nº 30. American Society of
1017 *Agronomy*, Crop Science Society of America, Soil Science Society of America.

1018 Soil Survey Staff, 2014. *Keys to Soil Taxonomy* (12th edition). United States Department of
1019 *Agriculture – Natural Resources Conservation Service*. 359 pp.

1020 Stoate, C., Boatman, N.D., Borralho, R.J., Carvalho, C.R., de Snoo, G.R., Eden, P., 2001.
1021 Ecological impacts of arable intensification in Europe. *J. Environ. Manage.* 63, 337–
1022 365. doi:10.1006/jema.2001.0473

1023 Stockle, C.O., 2001. Environmental impact of irrigation: A review. *Department of Biological*
1024 *Systems Engineering Newsletter*, Washington State University.

1025 Sutton, M.A., Howard, C.M., Erismann, J.W., Billen, G., Bleeker, A., Grennfelt, P., van Grisven,
1026 H., Grizzetti, B., 2011. *The European Nitrogen Assessment: Sources, Effects and*
1027 *Policy Perspectives*. Cambridge University Press, Cambridge, UK.

1028 Thorburn, P.J., Biggs, J.S., Attard, S.J., Kemei, J., 2011. Environmental impacts of irrigated
1029 sugarcane production: Nitrogen lost through runoff and leaching. *Agric. Ecosyst.*
1030 *Environ.* 144, 1–12. doi:10.1016/j.agee.2011.08.003

1031 USEPA, 2015. *Connectivity of Streams & Wetlands to Downstream Waters: A Review &*
1032 *Synthesis of the Scientific Evidence*. EPA/600/R-14/475F. Available at:
1033 <https://archive.epa.gov/> [Accessed April 2017]

- 1034 Villalobos, F.J., Mateos, L., Quemada, M., Delgado, A., Fereres, E., 2016. Control of Salinity, in
1035 Villalobos, F.J., Fereres, E. (Eds.), Principles of Agronomy for Sustainable Agriculture.
1036 Springer, Cham, Switzerland. doi:10.1007/978-3-319-46116-8
- 1037 Wang, R., Kang, Y., Wan, S., Hu, W., Liu, S., Jiang, S., Liu, S., 2012. Influence of different
1038 amounts of irrigation water on salt leaching and cotton growth under drip irrigation in an
1039 arid and saline area. *Agric. Water Manag.* 110, 109–117.
1040 doi:10.1016/j.agwat.2012.04.005
- 1041 Withers, P.J., Sharpley, A.N., 1995. Phosphorus fertilisers, in: Rehgigl, J.E. (Ed.), *Soil*
1042 *Amendments and Environmental Quality*. CRC Press, Inc., Boca Raton, Florida, USA,
1043 pp. 65–107.
- 1044
- 1045
- 1046

1047 Table 1. Crops cultivated and summary of agronomic management in Landazuria for
 1048 the period 2007-2016. Source: CR-EI Ferial (2017), INTIA (2017) and Government of
 1049 Navarre (2017).

	07	08	09	10	11	12	13	14	15	16	Average
Crops (%)¹											
Irrigated (252.9 ha)											
Maize	43	43	45	22	36	35	61	61	50	44	46
Winter cereals	10	34	24	14	19	31	30	30	42	24	27
Tomato	7	6	9	17	25	13	0	1	2	2	8
Onion	17	9	9	21	13	0	0	0	0	0	7
Others	31	18	23	39	27	41	26	15	38	47	22
Double cropping ²	8	10	10	13	20	20	17	7	32	17	10
N-fertilization (kg ha ⁻¹ year ⁻¹)	201	204	203	166	224	223	236	n.a.	n.a.	n.a.	209
Rainfed (170.5 ha)											
Fallow land	62	40	55	32	64	35	65	32	73	32	49
Witer cereals	36	56	43	64	34	63	33	66	26	66	49
Others	2	4	2	4	2	2	2	2	1	2	2
N-fertilization (kg ha ⁻¹ year ⁻¹)	28	28	33	41	27	25	42	n.a.	n.a.	n.a.	32

1050

1051

¹ 2007-2013: face-to-face inquiries, Life-Nitrates Project; 2014-2016: Estimated from Common Agricultural Policy

1052

declarations and Irrigation Authority (CR-EI Ferial) information.

1053

² Surface with two different crops in a single agricultural year (the sum of crops surfaces exceeds 100%).

1054

n.a.: not available.

1055

1056

1057 Table 2. Some characteristics of precipitation, irrigation (estimated) and discharge recorded at
 1058 Landazuria. Runoff coefficient = Discharge / (Precipitation + Irrigation).
 1059

	Precipitation [mm]		Irrigation [mm]		Discharge [mm]		Runoff coefficient [%]	
	Mean	S.D.	Mean	S.D.	Mean	S.D.	Mean	S.D.
Autumn	126	50	8	3	27	10	21.3	5.9
Winter	109	48	12	8	22	10	18.0	5.4
Spring	127	40	106	24	24	13	9.9	4.9
Summer	75	41	210	17	29	9	10.0	2.2
Annual	437	92	337	34	102	30	12.9	2.8

1060

1061

S.D.: Standard deviation.

1062

1063

1064 Table 3. Values of discharge, nitrate (NO₃-), phosphate (PO₄³⁻), sediment and total dissolved
 1065 solids (TDS) yields for the hydrological years. Estimated values representative of the whole
 1066 study period are presented in the line 2007-2016.

Agric. year (Sep-Aug)	Discharge mm	NO ₃ -N yield kg ha ⁻¹	PO ₄ ³⁻ -P yield g ha ⁻¹	Sediment yield kg ha ⁻¹	TDS yield kg ha ⁻¹
2007	143	n.a.	n.a.	n.a.	n.a.
2008	83	n.a.	n.a.	n.a.	n.a.
2009	98	39.0	n.a.	94	2,601
2010	88	n.a.	n.a.	n.a.	n.a.
2011	87	38.3	14	54	1,828
2012	48	22.2	13	76	996
2013	138	63.6	37	159	2,904
2014	88	33.9	52	1154	1,703
2015	147	45.2	97	740	2,640
2016	96	29.3	24	42	1,711
Average (S.D.)	102	38.8 (13.2)	39 (32)	360 (473)	1,798 (680)

1067

S.D.: Standard deviation

1068

n.a.: not available (not enough reliable data for an adequate estimation).

1069

1070

1071 Table 4. Correlation matrix among selected variables using the non-parametric Spearman's rho
 1072 (Helsel and Hirsch, 2002). Only significant correlations ($p < 0.05$) are presented. The case of the
 1073 Spearman's rho coefficient is selected according with the strength of the correlation (absolute
 1074 values of rho: < 0.40 ; $0.40-0.70$; > 0.70).


1075

	Q	NO ₃ ⁻	PO ₄ ³⁻	Sed.	TDS	N-L	P-L	Sed.-L	TDS-L
Q									
NO ₃ ⁻	-0.18								
PO ₄ ³⁻		-0.22							
Sed.	+0.19	-0.13	+0.34						
TDS		+0.41	-0.39	-0.17					
N-L	+0.92	+0.18	-0.11	+0.12					
P-L	+0.46	-0.29	+0.83	+0.39	-0.39	+0.34			
Sed-L	+0.54	-0.17	+0.28	+0.91	-0.18	+0.45	+0.53		
TDS-L	+0.94		-0.12	+0.14	+0.22	+0.93	+0.34	+0.47	

1076

1077

1078




1080 Figure 1. A) Network of experimental watersheds in Navarre (Spain). Main rivers (blue
 1081 lines) and irrigated surface (green). B) Agricultural plots in Landazuria. C) Soil map
 1082 following Soil Survey Staff (2014). Source: Government of Navarre (2005) and IDENA
 1083 (2017).

1084

1085

1086

1087



1088

1089 Figure 2. Relationship between chloride (Cl⁻) and total dissolved solids (TDS) concentrations in


1090 Landazuria outlet.

1091

1092

1093


1094



1095

1096 Figure 3. Monthly average discharge, precipitation and irrigation. Vertical bars indicate standard
1097 deviation.


1098




1099

1100 Figure 4. Selected two-month periods during the non-irrigated (A) and the irrigated (B) seasons
1101 of the year 2016 in which the influence of daily irrigation schedules was clearly observed.


1102



1103 Figure 5. Nitrate concentration (NO_3^- ; 95%-confidence interval on the monthly median) and daily
 1104 average discharge in Landazuria outlet.
 1105



1106
 1107 Figure 6. Phosphate concentration (PO_4^{3-} ; 95%-confidence interval on the monthly median) and
 1108 daily average discharge in Landazuria outlet.
 1109




1110

1111 Figure 7. Sediment concentration (95%-confidence interval on the monthly median) and daily
 1112 average discharge in Landazuria outlet.

1113

1114



1115

1116 Figure 8. Total suspended solids concentration (TDS; 95%-confidence interval on the monthly
 1117 median) and daily average discharge in Landazuria outlet.

1118