

## Male Batterers with and without Psychopathy: An Exploratory Study in Spanish Prisons

Running head: Psychopathy among batterers

Enrique Echeburúa<sup>1</sup>  
Javier Fernández-Montalvo<sup>2</sup>

- (1) Department of Psychology  
University of the Basque Country  
Avda. de Tolosa, 70  
20018 San Sebastián (Spain)  
Phone: (+34) 943 01 83 29  
E-mail: [enrique.echeburua@ehu.es](mailto:enrique.echeburua@ehu.es)
- (2) Department of Psychology and Education  
Public University of Navarre  
Campus of Arrosadía  
31006 Pamplona (Spain)  
Phone: (+34) 948 16 98 30  
E-mail: [fernandez.montalvo@unavarra.es](mailto:fernandez.montalvo@unavarra.es)

*Authors' Note:* This research was supported by an agreement between the University of the Basque Country and the Correctional Institution of Spain. Please address correspondence to Enrique Echeburúa, Department of Psychology, University of the Basque Country, Avda. de Tolosa, 70. 20018 San Sebastián (Spain). E-mail: [enrique.echeburua@ehu.es](mailto:enrique.echeburua@ehu.es)

**ABSTRACT**

The aim of this study was to analyze the differential profile of male batterers in prison with and without psychopathy. The sample consisted of 162 perpetrators sentenced for a serious offence against their intimate partner. The prevalence of psychopathy or psychopathic traits, established according to the PCL-R, was about 12%. The psychopathic batterers were younger, more impulsive and suspicious, less empathetic and with lower self-esteem than non-psychopathic batterer. However, the psychopathic batterers were not engaged in intimate femicide more often than the non-psychopathic batterers. The severity of the crime was similar in both groups, so it may be attributed to other variables (e.g., substance abuse, intoxication at the time of the offense, violence history, jealousy, etc.). Implications of these results for further research and clinical practice are commented on.

**Keywords:** Male batterers; psychopathy; prison; femicide.

## INTRODUCTION

There is an explosive growth of family violence research, but little is yet known about batterers in prison. Severely violent men have been found to have lower socioeconomic status, more disrupted attachment patterns and greater frequency of witnessing violence in their family of origin than all other groups of men (Schumacher, Felbau-Kohn, Smith & Heyman, 2001; Sugarman & Hotaling, 1989). But the discontinuities in the literature are often a consequence of the failure to recognize the heterogeneity among aggressors (O'Leary, 1993).

Developing reliable and valid classification systems would help identify different types of men who batter on the basis of both violent behavior and psychological characteristics. Such typologies could help identify different processes leading to gender violence (what is important for the development of one type of batterer may be irrelevant for the development of other types) and could also support the development of specific treatment practices for different male batterers (Aldarondo, 1998).

According to Holtzworth-Munroe & Stuart (1994), there are three main types of male batterers: family-only, dysphoric/borderline and generally violent/antisocial. Antisocial batterers are estimated to constitute approximately 25% of batterer samples, or even more if data come from treatment programs with court-mandated batterers (Hart, Dutton & Newlove, 1993). These men engage in violence outside their intimate relationship, have extensive criminal records and are usually affected by mental disorders (substance abuse problems, psychopathy, etc.). Moreover they tend to have hostile attitudes towards women, low empathy, and have the highest rate of alcohol dependence and previous convictions

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

(Hamberger & Hastings, 1988; Huss & Langhinrichsen-Rohling, 2000; Quinsey, Harris, Rice & Cormier, 1998; White & Gondolf, 2000).

There is a great deal of concern about identifying batterers in prison with mental disorders (Echeburúa, Fernández-Montalvo & Amor, 2003; Fernández-Montalvo, Echeburúa & Amor, 2005). The vast majority of jail inmates have substance use disorders and/or personality disorders (Nicholls, Roesch, Olley, Ogloff & Hemphill, 2005).

Regarding the personality disorders, the importance of violence in psychopathic symptomatology has always been clear, and is well represented in current diagnostic criteria: those for antisocial personality disorder in the *DSM-IV-TR* (American Psychiatric Association, 2000); those for dissocial personality in the *ICD-10* (World Health Organization, 1990); and those for psychopathy in the *Psychopathy Checklist-Revised (PCL-R)* (Hare, 1991). Each set contains one criterion directly related to a history of irritability, hostility, and aggression, including overt physical violence. In addition, each set contains several criteria that are indirectly related to aggression or violence (e.g., callousness, lack of remorse) (Hare, 2001).

Personality disorders have frequently been identified among perpetrators of domestic violence, the most commonly diagnosed being antisocial, borderline, and narcissistic (Hamberger & Hastings, 1988, 1991; Huss & Langhinrichsen-Rohling, 2000). That is, the men who batter their female partners are a heterogeneous group of individuals, differing in areas such as severity of abuse, generality of aggression and psychopathology.

If psychopaths represent a major challenge for the criminal justice system in democratic societies (Lösel, 2001), it is necessary to have more empirical knowledge about this subgroup of batterer men and to design

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

prediction instruments, such as *Danger Assessment* (Campbell, 1995).

Only an accurate understanding of the variability within batterer men will let us achieve the development of sound assessment instruments and effective treatments (Aldarondo, 1998; White & Gondolf, 2000).

The aim of this study is analyze the differential profile of male batterers in prison with and without psychopathy in order to design specific intervention programs for these types of perpetrators at a later stage. This purpose is relevant because of the scarcity of studies about this topic and because current treatments are less successful with this kind of perpetrators (Dutton, 2003). As a main hypothesis, psychopathic batterers would be expected to be involved in a high prevalence rate of intimate femicide, to have a previous history of psychiatric problems, to be affected by a low self-esteem and a high impulsivity and to hold more cognitive distortions (e.g., attitudes more supportive of domestic violence and sympathetic to male batterers) and psychopathological symptoms than non-psychopathic batterer.

## **METHOD**

### **Participants**

The sample for this study consisted of 162 participants sentenced for a serious offence of violence against their intimate partner. These participants are part of ongoing research on the effectiveness of a pilot program of psychological intervention with prison inmates convicted of violence against women that is currently running in eighteen Spanish prisons (2005 and 2006). All the offenders were incarcerated in national prisons (all jails belong to this system in Spain) in medium-security wings and gave informed consent to participate in the study.

Those selected for the sample were required to be: a) adult males

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

(between 18 and 65 years old); and b) serving a sentence for a serious offence in relation to gender violence against their partner. All participants took part voluntarily in the program, having been properly informed of its characteristics, and being allowed to withdraw from the study without penalty.

### **Assessment measures**

#### *a) Psychopathy*

The *Psychopathy Checklist-Revised (PCL-R)* (Hare, 1991; Spanish version by Moltó & Torrubia, 2000) is a semi-structured interview and a set of ratings based on the interview and corroboration from multiple information sources (case story reviews, interviews with family members, criminal and psychiatric records). This instrument could not provide a valid assessment of psychopathy in the absence of this additional corroboratory information. Specific scoring criteria are used to rate each of 20 items on a 3-point scale (0,1,2) according to the extent to which it applies to a given individual. Total scores can range from 0 to 40 and reflect the degree to which the individual matches the prototypical psychopath. The mean score is about 18-20 (SD about 7-8) in forensic psychiatric populations. A score of 30 typically is used as a diagnostic cut-off for psychopathy, but it is possible to adopt less stringent cut-offs to evaluate psychopathic tendencies (20 or more points). Findings from several studies attest to this instrument's predictive validity (e.g., Serin & Amos, 1995).

#### *b) Cognitive and Empathic Variables*

The *Inventory of Distorted Thoughts about Women* (Echeburúa & Fernández-Montalvo, 1998) comprises a checklist of 13 binary items aimed at detecting irrational thoughts in the aggressor that are related to sexual roles and the inferiority of women. Each affirmative response scores 1 point, so that

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

the inventory score ranges between 0 and 13 points. The higher the score, the greater the number of women-related cognitive distortions.

The *Inventory of Distorted Thoughts on the Use of Violence* (Echeburúa & Fernández-Montalvo, 1998) comprises a checklist of 16 binary items aimed at detecting irrational thoughts in the aggressor that are related to the use of violence as an acceptable way of resolving conflicts. Each affirmative response scores 1 point, so that the inventory score ranges between 0 and 16 points. The higher the score, the greater the number of cognitive distortions connected with the use of violence as an acceptable way of resolving conflicts.

The *Interpersonal Response Index* (Davis, 1980; Spanish version by Garrido & Beneyto, 1995) consists of 28 items that assess four components of empathy: *fantasy* (capacity for imagination and identification with fictional characters), *awareness of perspective* (capacity to appreciate the point of view of others), *empathic interest* (capacity for showing concern for persons who have negative experiences), and *personal grief* (capacity to feel the negative emotions of others as one's own). Each of the 28 items is marked on a Likert-type scale that ranges from 0 (*absolute disagreement*) to 4 (*absolute agreement*). The full range of the scale is, therefore, from 0 to 112. The higher the score, the greater the empathic capacity.

### c) *Psychopathological and Personality Variables*

The *Symptom Checklist-90-Revised* (*SCL-90-R*; Derogatis, 1975; Spanish version by González de Rivera, 2002) is a self-administered general psychopathological assessment questionnaire. It comprises 90 items with five alternatives for each item on a Likert-type scale, ranging from 0 (*none*) to 4 (*very much*). As it has been shown to be sensitive to therapeutic change, it may be used for either single or repeated assessments (Echeburúa,

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

Fernández-Montalvo & Amor, 2006). The *SCL-90-R* consists of nine areas of primary symptoms (somatization, obsessive-compulsive, interpersonal sensitivity, depression, anxiety, hostility, phobic anxiety, paranoid ideation and psychoticism). It also provides three overall indices that reflect the participant's overall level of severity.

The *State-Trait Anger Expression Inventory (STAXI)* (Spielberger, 1988; Spanish version by Miguel-Tobal, Casado & Cano-Vindel, 2001) consists of 15 items related to state-anger (the intensity of the emotion of anger in a specific situation) and a further 10 items related to trait-anger (the individual disposition to experience anger habitually). The range of scores is from 15 to 60 on the state-anger scale and from 10 to 40 on the trait-anger scale. The *STAXI* also has a third subscale of 24 items connected with the form of expressing anger (anger expression-out, anger expression-in, and anger control).

The *Impulsivity Scale (BIS-10)* (Barratt, 1985; Spanish version by Luengo, Carrillo de la Peña & Otero, 1991) consists of 33 items aimed at assessing how impulsive participants are. Scores from 0 to 4 on a Likert-type scale provide a total scale range of between 0 and 132. The higher the score, the stronger the presence of each trait measured on each subscale. The sum of all the subscales gives the total score.

The aim of the *Self-Esteem Scale* (Rosenberg, 1965; Spanish version by Fernández-Montalvo & Echeburúa, 1997) is to assess the feeling of satisfaction that a person has about him or herself. There are 10 general items, each carrying a score of between 1 and 4 on a Likert-type scale, giving a questionnaire range of 10 to 40. The higher the score, the greater the level of self-esteem. The cut-off point for the adult population is 29 points. Test-

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.



retest reliability is .85, and the internal consistency *alpha* coefficient is .92. Convergent validity and discriminant validity are likewise satisfactory (*cf.* Zubizarreta, Sarasua, Echeburúa, Corral, Saucá & Emparanza, 1994).

All the instruments above described have proven to have good psychometric characteristics for this specific population in Spanish prisons (*cf.* Echeburúa *et al.*, 2003; Fernández-Montalvo *et al.*, 2005).

### **Procedure**

All the participants completed the questionnaires individually in the psychologist's presence during pretreatment assessment before the intervention program. This assessment was carried out during April and May 2005 by correctional psychologists under the direction of the authors of this study.

This is a transversal study that forms part of a wider study looking at a penitentiary psychological treatment.

### **RESULTS**

In this study the level of statistical significance chosen for comparison between groups was  $p < .05$ .

#### **Psychopathy prevalence**

In terms of psychopathy, according to the results of the *PCL-R*, there were 20 people (the 12% of the sample) who met the criteria for psychopathy (30 or more points) or probable psychopathy (psychopathic traits) (20 or more points).

#### **Comparison between male batterers with and without psychopathy**

Sociodemographic and penal characteristics and results of the comparison between male batterers with and without psychopathy are shown in Table 1. As can be seen, there is only one significant difference, related to

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

age, with the psychopathic batterers being younger than those without psychopathy. Although in a non-significant trend, the psychopathic batterers also were more likely to have a previous history of psychiatric problems than were the batterers without psychopathy. The main disorders for which professional assistance had been given were addictive behaviors and depression.

There were not any differences in regard to intimate femicide. The femicide prevalence in the whole sample was about 17%. Femicide was not more prevalent in psychopathic batterers than in non-psychopathic batterers.

---

*TABLE 1*

---

With regard to the rest of studied variables, the level of empathy and self-esteem was significantly lower and the level of the impulsivity and of the interpersonal sensitivity (suspiciousness) was significantly higher in psychopathic batterers than in the rest of batterers (Tables 2 and 3).

---

*TABLES 2 and 3*

---

## **DISCUSSION**

This study deals with 162 men in eighteen prisons in Spain who were convicted of serious violence against women, including 27 men who actually murdered their intimate partners. It attempts to characterize the profile of batterer men according to the diagnosis of psychopathy.

Only a small group of the whole sample of batterer men (1 out of 8) in our study fits the profile of psychopathic batterer, that is, of a cold-blooded aggressor who, with no previous emotional instability, commits a serious

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

offence in an insensitive and cruel way. Indeed, most of the batterer men in prison are not psychopaths, but men who have hostile feelings against women, tend to be affectively instable, abuse of alcohol or drugs, are affected by a possible impulse control disorder or an intermittent explosive disorder, or have been engaged in a fit of rage or jealousy.

The prevalence of psychopathy or probable psychopathy (psychopathic traits) in batterer men incarcerated for a serious offence against their partner in our study is about 12%, somewhat lower than in general prison populations (about 15-25%) as indicated by other studies (Hare, 2001). But prison inmates who meet the criteria for psychopathy or who have a significant number of psychopathic traits have been found to be at much higher risk for recidivism and violence than are other prisoners (Grann & Wedin, 2002; Harris, Rice & Quinsey, 1993; Stadtland, Kleindienst, Kröner, Eidt & Nedopil, 2005; Walters, 2003). Thus, the risk for violence against women is increased by a diagnosis of psychopathy in the batterers, as also has been found in other studies (Danielson, Moffit, Caspi & Silva, 1998).

The psychopathic batterers in this study were younger, more impulsive and suspicious, less empathetic and with lower self-esteem than non-psychopathic batterers. However, they did not commit intimate femicide more often than the non-psychopathic batterers. The severity of the crime was similar in both groups, so it may be attributed to other variables (e.g., substance abuse, intoxication at the time of the offense, violence history, jealousy, etc.) elsewhere studied (Echeburúa *et al.*, 2003; Fernández-Montalvo *et al.*, 2005). Indeed, according to Campbell (1995), the main risk factors for intimate femicide include access to guns, threats with weapons, serious injury in prior abusive incidents, threats of suicide, drug

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

or alcohol abuse, forced sex of female partner and obsessiveness/extreme jealousy/extreme dominance.

As in other studies (Holtzworth-Munroe & Stuart, 1994; Huss & Langhinrichsen-Rohling, 2000), our study gives empirical support to the existence of a subgroup of batterers that can be characterized as exhibiting significant psychopathic traits. This subgroup of domestically violent men exhibit more antisocial behavior, express more generalized violence, and are more likely to abuse illicit substances. However, there is a lack of consistency between our results and the estimate of Holtzworth-Munroe & Stuart (1994) that about 25% of batterers belonging to the antisocial type. This discrepancy may be attributed to a more strict definition of psychopathy of *PCL-R* in comparison with antisocial personality.

There are some limitations in this study. One potential limitation in the field of domestic violence is that *PCL-R* was designed to predict general violence and criminal recidivism so it gathers little or no information on partner abuse (Dutton & Kropp, 2000). Also, this is an exploratory study with the sample size of psychopaths not being large enough to generate generalizable and reliable findings. Likewise, the weight of social desirability in the low level of psychopathology found, measured with self-reports, cannot be disregarded in this study. Such desirability in the group of batterers in prison may be greater than expected. In short, pretending a degree of normality in front of examiners may be one way of gaining faster access to probation (Fernández-Montalvo *et al.*, 2005). Further studies are needed to test these conclusions.

Future research should focus on replicating these data with larger samples and designing tailored programs for these kind of perpetrators.

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

Intervening with the psychopath batterers is relevant to practitioners and researchers in both the domestic violence and correctional psychology fields because current treatments are less successful with this kind of perpetrators (Dutton, 2003; Lösel, 2001).

## REFERENCES

- Aldarondo, E. (1998). Perpetrators of Domestic Violence. In N.N. Sing (ed.). *Applications in Diverse Populations (A.S. Bellack & M. Hersen. Comprehensive Clinical Psychology, vol. 9, pp. 437-452)*. Amsterdam. Elsevier Science.
- American Psychiatric Association (2000). *Diagnostic and Statistical Manual of Mental Disorders (4<sup>th</sup>. ed. rev.)*. Washington, DC. Author.
- Barratt, E.S. (1985). Impulsiveness subtraits: arousal and information procesing. In J.T. Spence & C.E. Itard (eds.). *Motivation, Emotion and Personality*. North Holland. Elsevier.
- Campbell, J.C. (1995). *Assessing Dangerousness: Violence by Sexual Offenders, Batterers, and Child Abusers*. Thousand Oaks, CA. Sage.
- Danielson, K.K., Moffit, T.E., Caspi, A. & Silva, P.A. (1998). Comorbidity between abuse of an adult and DSM-III-R mental disorders: Evidence from an epidemiological study. *American Journal of Psychiatry, 155*, 131-133.
- Davis, M.H. (1980). *Interpersonal Reactivity Index. A multidimensional approach to individual differences in empathy*. Washington. American Psychological Association.
- Derogatis, L.R. (1975). *The SCL-90-R*. Baltimore. Clinical Psychometric Research.
- Dutton, D. (2003). Theoretical approaches to the treatment of intimate violence perpetrators. *Journal of Aggression, Maltreatment and Trauma, 7*, 7-23.
- Dutton, D.G. & Kropp, P.R. (2000). A review of domestic violence risk instruments. *Trauma, Violence and Abuse, 1*, 171-182.
- Echeburúa, E. & Fernández-Montalvo, J. (1998). Hombres maltratadores (*Male batterers*). In E. Echeburúa & P. Corral. *Manual de violencia familiar (Handbook of Family Violence, pp. 70-175)*. Madrid. Siglo XXI.
- Echeburúa, E., Fernández-Montalvo, J. & Amor, P.J. (2003). Psychopathological profile of men convicted of gender violence: A study in the prisons of Spain. *Journal of Interpersonal Violence, 18*, 798-812.
- Echeburúa, E., Fernández-Montalvo, J. y Amor, P.J. (2006). Psychological treatment of men convicted of gender violence: A pilot-study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology, 50*, 57-70.
- Fernández-Montalvo, J. & Echeburúa, E. (1997). *Manual práctico del juego patológico. Ayuda para el paciente y guía para el terapeuta (Handbook of*
- Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology, 51* (3), 254-263.

*pathological gambling: Guide to the therapist and self-help booklet to the patient*). Madrid. Pirámide.

Fernández-Montalvo, J., Echeburúa, E. & Amor, P.J. (2005). Aggressors against women in prison and in community: An exploratory study of a differential profile. *International Journal of Offender Therapy and Comparative Criminology*, 49, 158-167.

Garrido, V. & Beneyto, M.J. (1995). *Control de la agresión sexual (Control of sexual aggression)*. Valencia. Cristóbal Serrano Villalba.

González de Rivera, J.L. (2002). *Versión española del SCL-90-R (Spanish version of SCL-90-R)*. Madrid. TEA.

Grann, M. & Wedin, I. (2002). Risk factors for recidivism among spousal assault and spousal homicide offenders. *Psychology, Crime and Law*, 8, 5-23.

Hamberger, L.K. & Hastings, J.E. (1988). Characteristics of male spouse abusers consistent with personality disorders. *Hospital and Community Psychiatry*, 39, 763-770.

Hamberger, L.K. & Hastings, J.E. (1991). Personality correlates of men who batter and non-violent men: Some continuities and discontinuities. *Journal of Family Violence*, 6, 131-148.

Hare, R.D. (1991). *The Hare Psychopathy Checklist-Revised*. Toronto. Multi-Health Systems.

Hare, R.D. (2001). Psychopaths and their nature. In A. Raine & J. Sanmartín (eds.). *Violence and psychopathy* (pp. 5-34). New York. Kluwer Academic/Plenum Publishers.

Harris, G.T., Rice, M.E. & Quinsey, V.L. (1993). Violence and recidivism of mentally disordered offenders: The development of a statistical prediction instrument. *Criminal Justice and Behavior*, 20, 315-335.

Hart, S.D., Dutton, D.G. & Newlove, T. (1993). The prevalence of personality disorder amongst wife assaulters. *Journal of Personality Disorders*, 7, 329-341.

Holtzworth-Munroe, A. & Stuart, G.L. (1994). Typologies of male batterers: Three subtypes and the differences among them. *Psychological Bulletin*, 116, 476-497.

Huss, M.T. & Langhinrichsen-Rohling, J. (2000). Identification of the psychopathic batterer: The clinical, legal, and policy implications. *Aggression and Violent Behavior*, 5, 403-422.

Lösel, F. (2001). Is effective treatment of psychopathy possible?. In A. Raine & J. Sanmartín (eds.). *Violence and psychopathy* (pp. 171-195). New York. Kluwer Academic/Plenum Publishers.

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

Luengo, M.A., Carrillo de la Peña, M.T. & Otero, J.M. (1991). The components of impulsiveness: a comparison of the I.7 Impulsiveness Questionnaire and the Barratt Impulsiveness Scale. *Personality and Individual Differences*, 12, 657-667.

Miguel-Tobal, J.J., Casado, M.I., Cano-Vindel, A. & Spielberger, C.D. (2001). *Versión española del STAXI-2 (Spanish version of STAXI-2)*. Madrid. TEA.

Moltó, J. & Torrubia, R. (2000). Standardization of the Hare Psychopathy Checklist-Revised in a Spanish Prison Sample. *Journal of Personality Disorders*, 14, 84-96.

Nicholls, T.L., Roesch, R., Olley, M.C., Ogloff, J.R. & Hemphill, J.F. (2005). *Jail Screening Assessment Tool (JSAT): Guidelines for Mental Health Screening in Jails*. Vancouver. Mental Health, Law, and Policy Institute.

O'Leary, K.D. (1993). Controversies regarding psychological explanations of family violence. In R.G. Gelles & D. Loseke (eds.). *Current controversies on family violence* (pp. 7-30). Newbury Park, CA. Sage.

Quinsey, V.L., Harris, G.T., Rice, M.E. & Cormier, C.A. (1998). *Violent offenders: Appraising and managing risk*. Washington, DC. American Psychological Association.

Rosenberg, A. (1965). *Society and the adolescent self-image*. New Jersey. Princenton.

Schumacher, J.A., Felbau-Kohn, S., Smith, A. & Heyman, R.E. (2001). Risk factors to male-to-female partner physical abuse. *Aggression and Violent Behavior*, 6, 281-352.

Serin, R.C. & Amos, N.L. (1995). The role of psychopathy in the assessment of dangerousness. *International Journal of Law and Psychiatry*, 18, 231-238.

Spielberger, C.D. (1988). *State-Trait Anger Expression Inventory*. Orlando, FL. Psychological Assessment Resources.

Stadtland, C., Kleindienst, N., Kröner, C., Eidt, M. & Nedopil, N. (2005). Psychopathic Traits and Risk of Criminal Recidivism in Offenders with and without Mental Disorders. *International Journal of Forensic Mental Health*, 4, 89-97.

Sugarman, D.B. & Hotaling, G.T. (1989). Violent men in intimate relationships : An analysis of risk markers. *Journal of Applied Social Psychology*, 19, 1034-1048.

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.



Walters, G.D. (2003). Predicting institutional adjustment and recidivism with the psychopathy checklist factor scores : A meta-analysis. *Law and Human Behavior, 27*, 541-558.

White, R.J. & Gondolf, E.W. (2000). Implications of personality profiles for batterer treatment. *Journal of Interpersonal Violence, 15*, 467-488.

World Health Organization (1990). *International Classification of Diseases and Related Health Problems (10th ed.)*. Geneva (Switzerland). Author.

Zubizarreta, I., Sarasua, B., Echeburúa, E., Corral, P., Saucá, D. & Emparanza, I. (1994). Consecuencias psicológicas del maltrato doméstico (Psychological consequences of family violence). In E. Echeburúa (ed.). *Personalidades violentas (Violent personalities, pp. 129-152)*. Madrid. Pirámide.

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology, 51* (3), 254-263.

**TABLE 1**  
**COMPARISON BETWEEN PARTICIPANTS WITH AND WITHOUT**  
**PSYCHOPATHY IN SOCIODEMOGRAPHICAL AND PENAL VARIABLES**

	Psychopaths N=20 ----- Mean (SD)	No psychopaths N=142 ----- Mean (SD)	<i>t</i>
Age	35.7 (9.2)	40.6 (9.6)	2.01 (p=.03)
Months in prison	30.7 (42.8)	20.5 (23.2)	1.5 (p=.10)
	Psychopaths N=20 ----- N (%)	No psychopaths N=142 ----- N (%)	$\chi^2$
<b>Marital status</b>			
Married	1 (5%)	28 (19.7%)	6,1 (p=.10)
Single	9 (45%)	31 (21.8%)	
Divorced	9 (45%)	75 (52.8%)	
Widowed	1 (5%)	8 (5.6%)	
<b>Education</b>			
None	2 (10%)	12 (8.4%)	.35 (p=.94)
Primary studies	15 (75%)	110 (77.4%)	
Secondary studies	2 (10%)	16 (11.3%)	
University	1 (5%)	4 (2.8%)	
<b>Socioeconomic status</b>			
Low	4 (20%)	29 (20.4%)	1.1 (p=.77)
Middle-low	10 (50%)	59 (41.5%)	
Middle	6 (30%)	49 (34.5%)	
Middle-high	0	5 (3.5%)	
<b>Previous psychiatric history</b>			
Yes	13 (65%)	58 (40.8%)	3.2 (p=.07)
No	7 (35%)	84 (59.2%)	
<b>Criminal records</b>			
Yes	8 (40%)	47 (33.1%)	.12 (p=.72)
No	12 (60%)	95 (66.9%)	
<b>Homicide</b>			
Yes	3 (15%)	25 (17.6%)	.08 (p=.97)
No	17 (85%)	117 (82.4%)	

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

**TABLE 2**  
**COGNITIVE DISTORTIONS AND PSYCHOPATHOLOGICAL AND**  
**PERSONALITY VARIABLES**

	Psychopaths N=20 ----- Mean (SD)	No psychopaths N=142 ----- Mean (SD)	<i>t</i>
Cognitive bias about women	3.5 (1.7)	4.1 (2.2)	1.06 (p=.24)
Cognitive bias about violence use	5.5 (2.4)	6.3 (2.3)	1.30 (p=.14)
Empathy	51.5 (12.7)	62.4 (14.6)	3.16 (p=.002)
State-anger	17.8 (3.5)	17.3 (5.3)	0.33 (p=.68)
Trait-anger	18.5 (6.7)	16.6 (4.9)	1.50 (p=.13)
Expression of anger	33.3 (15.5)	27.1 (17.1)	1.54 (p=.12)
Impulsivity	55.3 (19.3)	46.8 (16.4)	2.12 (p=.03)
Self-esteem	25.9 (5.1)	29.4 (5.1)	2.92 (p=.004)

Echeburúa, E. y Fernández-Montalvo, J. (2007). Male batterers with and without psychopathy: an exploratory study in Spanish prisons. *International Journal of Offender Therapy and Comparative Criminology*, 51 (3), 254-263.

**TABLE 3**  
**COMPARISONS IN THE SCL-90-R (Percentiles)**

	Psychopaths N=20		No psychopaths N=142		<i>t</i>
	Mean	(SD)	Mean	(SD)	
<b>Global Symptoms Index</b>	56.4	(38.9)	63.1	(29.2)	0.86 (p=.35)
<b>Positive Symptoms Distress Index</b>	40.2	(34.7)	50.4	(29.6)	1.31 (p=.15)
<b>Positive Symptoms Total</b>	78.4	(26.7)	67.5	(28.8)	1.48 (p=.11)
<b>Somatization</b>	74.1	(23.9)	62.8	(31.4)	1.54 (p=.12)
<b>Obsessive-compulsive</b>	63.6	(32.3)	56.3	(30.9)	0.98 (p=.32)
<b>Interpersonal sensibility</b>	75	(25.9)	59.6	(29.1)	2.24 (p=.02)
<b>Depression</b>	77.1	(22.4)	69.5	(25.1)	1.25 (p=.20)
<b>Anxiety</b>	63.7	(35.3)	57.2	(33.3)	0.80 (p=.41)
<b>Hostility</b>	50.9	(35.4)	37.6	(33.1)	1.67 (p=.10)
<b>Phobic anxiety</b>	51.9	(36.9)	46.3	(36.1)	0.63 (p=.51)
<b>Paranoid ideation</b>	72.5	(30.1)	62.5	(32.2)	1.29 (p=.19)
<b>Psychoticism</b>	74.5	(24.9)	62.7	(33.8)	1.49 (p=.13)