

Supporting Information

Baroja-Fernández et al. 10.1073/pnas.1117099109




Fig. S1. UDPG (Upper) and ADPG (Lower) detection in 30 μ L of SUS assay reaction mixture including recombinant SUS1. The SUS assay mixture contained 50 mM Hepes (pH 7.0), 200 mM sucrose, 2 mM of UDP or ADP, and recombinant SUS1 (10 ng). After 3 min at 37 $^{\circ}$ C, reactions were stopped by boiling the assay mixture for 1 min. UDPG and ADPG were then measured by HPLC as described in *Materials and Methods*. In *A*, recombinant SUS1 heated at 100 $^{\circ}$ C for 1 min was included in the assay mixture. In *B*, the assay mixture included nonheated, active recombinant SUS1. In *C* and *D*, once the SUS reaction was stopped, 5 nmol of commercially available UDPG/ADPG and 1 U of USPP/ASPP were added to the assay mixture, respectively. UDPG and ADPG elution peaks are indicated by arrows. UDP and ADP elution times were 40.3 min and 40.5 min, respectively.





Fig. 54. UDPG detection in 30 μ L of SUS assay reaction mixture including crude extracts from WT and *sus1/sus2/sus3/sus4* leaves. The SUS assay mixture contained 50 mM Hepes (pH 7.0), 200 mM sucrose, 2 mM UDP, and crude extract. After 3 min at 37 $^{\circ}$ C (still under initial velocity conditions, see Fig. S3) reactions were stopped by boiling the assay mixture for 1 min. UDPG was then measured by HPLC as described in *Materials and Methods*. In *A*, crude extract heated at 100 $^{\circ}$ C for 1 min was included in the assay mixture. In *B*, the assay mixture included fresh crude extract. In *C* and *D*, once the SUS reaction was stopped, 5 nmol of commercially available UDPG and 1 U of USPP were added to the assay mixture, respectively. UDPG elution time was 27.5 min. UDPG elution peaks are indicated by arrows.






Fig. S6. Stages to produce the constructs used in this work. Primers used for the synthesis and cloning of full-length SUS-encoding cDNAs into pDEST17 expression vector are listed in Table S2.

Table S1. Primers used to confirm homozygous *sus1/sus2/sus3/sus4* and *sus5/sus6* mutants (Fig. S5)

Gene	Orientation	Sequence
<i>SUS1</i>	LP	5'-CTCAAGAGTGCAAGGATCAGG-3'
	RP	5'-ACGCTGAACGTATGATAACGC-3'
<i>SUS2</i>	LP	5'-CATGGGTAATGGTTTGGTTG-3'
	RP	5'-ATGCGGAGACAAAATCACAAC-3'
<i>SUS3</i>	LP	5'-TTGGAGACCAGCGTCTGATAC-3'
	RP	5'-ATCGATGTGTTTGATCCGAAG-3'
<i>SUS4</i>	LP	5'-CAATAGGATCCAGAACCTAAAC-3'
	RP	5'-CCGTGAACCTACACGGTACAAC-3'
<i>SUS5</i>	LP	5'-ATTTCCTTTTACCGCACAAG-3'
	RP	5'-TTGTTTGGCCAGTTTCTGATC-3'
<i>SUS6</i>	LP	5'-TGACACGGTTAATACCGGAAG-3'
	RP	5'-ATCCATCTGAATTCCTTCTG-3'

LP, left primer; RP, right primer.

Table S2. Primers used for the synthesis and cloning of full-length *SUS* encoding cDNAs into pDEST17 expression vector

Primer	Orientation	Sequence
<i>SUS1</i> -attB1	Forward	5'- ggggacaagttgtacaaaaaagcaggctta atggcaaacgctgaacgtatg-3'
<i>SUS1</i> -attB2	Reverse	5'- ggggaccactttgtacaagaaagctgggtat caatcatctgtgcaagagg-3'
<i>SUS3</i> -attB1	Forward	5'- ggggacaagttgtacaaaaaagcaggctta atggcaaaccttaagctcac-3'
<i>SUS3</i> -attB2	Reverse	5'- ggggaccactttgtacaagaaagctgggtat cagtcacgcggtgaagg-3'

Primer sequences for attB sites (Fig. S6) are indicated in bold.

Table S3. Primers used in real time PCR analyses of *SUS* expression

Gene	Orientation	Sequence
<i>18S RNA</i> (At3g41768)	Forward	5'-GGGCATTGCTATTTTCATAGTCAGAG-3'
	Reverse	5'-CGGTTCTTGATTAATGAAAACATCCT-3'
<i>SUS1</i> (At5g20830)	Forward	5'-AGTTCACTGCGGATATTTTCGC-3'
	Reverse	5'-CCCAACAGTTTCTTTGCTTCCA-3'
<i>SUS2</i> (At5g49190)	Forward	5'-TGCCATGAATAATGCCGATTTC-3'
	Reverse	5'-TTGCCCAACATTGTTCTTGCTT-3'
<i>SUS3</i> (At4g02280)	Forward	5'-GACCAGACTGATGAGCATGTCG-3'
	Reverse	5'-TCTTCACTTTGTCGAGCCTCG-3'
<i>SUS4</i> (At3g43190)	Forward	5'-AAGGAATCGTTCGCAAATGG-3'
	Reverse	5'-TTTCAGCGGCAACATCCTC-3'
<i>SUS5</i> (At5g37180)	Forward	5'-GAACGTAGCGTTCGAGTAA-3'
	Reverse	5'-TGAGATATTGCGTGCTCGAT-3'
<i>SUS6</i> (At1g73370)	Forward	5'-CGGAGGCCAGGTTGTTTACAT-3'
	Reverse	5'-AGGCTTGAATCCGAGACCTTGT-3'