

# Asymmetric Assembly of All-Carbon Tertiary/Quaternary Nonadjacent Stereocenters Through Organocatalytic Conjugate Addition of $\alpha$ -Cyanoacetates to a Methacrylate Equivalent


Igor Iriarte,<sup>[a]</sup> Silvia Vera,<sup>[a]</sup> Eider Badiola,<sup>[a]</sup> Antonia Mielgo,<sup>[a]</sup> Mikel Oiarbide,<sup>[a]\*</sup> Jesús M. García,<sup>[b]</sup> José M. Odriozola,<sup>[b]</sup> and Claudio Palomo<sup>\*[a]</sup>

Dedication ((optional))


**Abstract:** An efficient highly diastereo- and enantioselective assembly of acyclic carbonyl fragments possessing nonadjacent all-carbon tertiary/quaternary stereoarrays is reported based on a Brønsted base-catalyzed Michael addition/ $\alpha$ -protonation sequence involving  $\alpha$ -cyanoacetates and 2,4-dimethyl-4-hydroxypenten-3-one as novel methacrylate equivalent.

## Introduction


Acyclic carbonyl compounds possessing multiple stereocenters are important building-blocks for the construction of complex natural products and bioactive molecules. Huge advances have been made to synthesize stereoselectively some of these stereoarrays, particularly carbonyl compounds with contiguous stereocenters at  $\alpha,\beta$ - or  $\beta,\gamma$ -positions (**A** and **B**, Fig. 1). In contrast, direct asymmetric entries to the  $\alpha,\gamma$ -branched analogs **C**, bearing two nonadjacent stereocenters, are less common,<sup>[1]</sup> and rarely involve construction of a quaternary stereocenter as in **D**.<sup>[2,3]</sup> Among the few precedents, Deng reported a Brønsted base-catalyzed Michael/ $\alpha$ -protonation reaction cascade which




**Figure 1.** Acyclic carbonyl compounds with different stereoarrays.



a) organocatalysis



b)  $\text{Ca}(\text{BOX})_2$  catalysis (Kobayashi 2008; addition of glycine Schiff bases; tertiary stereocenters only)



**Figure 2.** Advances in tandem addition/protonation approaches for the asymmetric assembly of  $\alpha,\gamma$ -nonadjacent stereocenters.

implies  $\alpha$ -chloroacrylonitrile as the Michael acceptor,<sup>[4]</sup> and later on the group of Chen and Xiao described<sup>[5]</sup> a similar tandem reaction involving ethyl 2-phthalimidoacrylate or ethyl  $\alpha$ -phosphonoacrylates as the doubly activated Michael acceptor (Fig. 2). However, the extension of this methodology to inherently less reactive  $\alpha$ -alkyl substituted Michael acceptors, i.e. methacrylates, remains challenging, despite the fact that the resulting  $\alpha$ -alkyl –and more specifically  $\alpha$ -methyl– carbonyl units are present in a number of natural products and bioactive targets. In that respect, Kobayashi has reported<sup>[6]</sup> a  $\text{Ca}(\text{BOX})_2$ -catalyzed conjugate addition of glycine Schiff bases to  $\alpha$ -substituted acrylate derivatives, albeit no example involving generation of a quaternary center was gathered. More recently Pihko has described<sup>[7]</sup> enantioselective Mukaiyama-Michael addition reactions of methacrolein via iminium activation that, however, led to an approximately 1:1 mixture of the two possible diastereomers. To the best of our knowledge, highly enantioselective Michael reactions with methacrylates or equivalents to provide carbonyl compounds with all-carbon tertiary/quaternary nonadjacent stereocenters have not been realized yet. Here we present an effective asymmetric direct entry to such stereoarrays that is founded upon the development of a new designed methacrylate equivalent in combination with Brønsted base catalysis.


[a] Prof. C. Palomo, Prof. M. Oiarbide, Dr. A. Mielgo, Dr. S. Vera, E. Badiola, I. Iriarte  
Departamento de Química Orgánica I  
Universidad del País Vasco UPV/EHU  
Manuel Lardizabal 3, 20018 San Sebastián, Spain  
Fax: (+34) 943015270  
E-mail: claudio.palomo@ehu.es

[b] Prof. J. M. García, Dr. J. M. Odriozola  
Departamento de Química Aplicada  
Universidad Pública de Navarra  
31006-Pamplona, Spain

Supporting information for this article is given via a link at the end of the document.


## Results and Discussion

**Hypothesis and working plan.** The challenge posed by  $\alpha$ -alkyl substituted Michael acceptors is not only associated to their attenuated reactivity against neutral C-pronucleophiles, but also concerns stereocontrol during the key C-C bond formation (step 1, Fig. 3a) and the subsequent  $\alpha$ -protonation (step 2).<sup>[8]</sup>




b) Working hypothesis:

- Step 1: precedents with  $\alpha$ -unsubstituted  $\alpha'$ -hydroxy enones
- Step 2: efficient control over enolate *E/Z* geometry



**Figure 3.** Construction of nonadjacent tertiary/quaternary stereocenters and working hypothesis for designed methacrylate equivalent.



Preliminary studies involving the reaction of 2-phenyl  $\alpha$ -cyanoacetate **2a**<sup>[9]</sup> and some elementary  $\alpha$ -substituted Michael acceptors revealed these difficulties. For instance (Scheme 1) attempts to react **2a** with methyl methacrylate in the presence of several mono- and bifunctional Brønsted base catalysts all led to recovery of unreacted material. Under similar conditions, 3-methylbutenone resulted essentially unreactive at ambient temperature; 60% conversion was hardly achieved only after 90 h at 50 °C. Finally, methacrolein was more reactive, but led to unselective reaction.<sup>[7]</sup>



**Scheme 1.** Difficulties in the addition of  $\alpha$ -cyanoester **2a** to  $\alpha$ -methyl  $\alpha,\beta$ -unsaturated ester, ketone or aldehyde under best reaction conditions.

Recently we have introduced  $\alpha$ -unsubstituted  $\alpha'$ -hydroxy enones as efficient acrylate equivalents in Brønsted base-catalyzed enantioselective conjugate additions.<sup>[10]</sup> Both the remarkable reaction acceleration and the high level of asymmetric induction observed in these reactions were rationalized by assuming H-

bond-mediated effective substrate-catalyst complexation, with the ketol moiety of substrate forming tight 1,4-proton chelates. Now we hypothesize that by using the parent  $\alpha$ -substituted  $\alpha'$ -hydroxy enones, the unique capacity of these type of bidentated substrates<sup>11</sup> to act as both H-bond donor and acceptor, in cooperation with a proper Brønsted base catalyst, may also be translated to the  $\alpha$ -stereocenter determining step 2. If so, a shortcut to the challenging construction of acyclic carbonyl adducts with nonadjacent stereocenters would be provided. Specifically, it is predicted that the evolved enolate from step 1 would preferentially adopt a *Z* configuration because of unfavourable  $A^{1,3}$  strain in the chelated *E* form (Fig. 3b), thus overcoming the problem of ill-defined enolate geometry in asymmetric  $\alpha$ -protonations.<sup>12, 13</sup> Eventually, the prevalence of such dynamic H-bond networks may also help during proton transfer (shuttle). To achieve high overall selectivity, however, both chiral units, namely the catalyst and the newly generated stereocenter at  $\gamma$ , have to work in concert in step 2, an inherent difficult of the process that at the outset remained unclear. In this respect, as far as we know,  $\alpha$ -substituted  $\alpha'$ -hydroxy enones have never been employed in catalytic asymmetric conjugate additions.<sup>14</sup>



**Scheme 2.** Reaction of **1** with  $\alpha$ -cyanoacetates catalyzed by chiral Brønsted bases.

**Catalysts screening and reaction optimization.** The investigation was started by studying the reaction of the newly prepared  $\alpha'$ -hydroxy enone **1**<sup>[15]</sup> with  $\alpha$ -substituted cyanoacetate **2a** in the presence of several bifunctional Brønsted bases.<sup>[16]</sup> Among the catalysts examined, i.e. (DHQD)<sub>2</sub>PYR **C1**, quinine **C2**, the thiourea-aminoquinine **C3**<sup>[17]</sup> and the squaramides **C4**,<sup>[18]</sup> **C5**<sup>[19]</sup> and **C6**,<sup>[20]</sup> the latter proved to be superior as shown in Table 1. Using this catalyst the reaction between 2-phenyl  $\alpha$ -cyanoacetate **2a** and **1** at ambient temperature afforded adduct **5a** with essentially perfect enantio- and diastereocontrol (dr >99:1, 99% ee; entry 6). This result indicates that not only the initial addition that renders the  $\gamma$ -stereocenter, but also the subsequent  $\alpha$ -protonation, both proceeded with remarkable face selectivity.

## FULL PAPER

Interestingly, this almost perfect chirality transfer was reproduced (entry 7) when the reaction was run at 50 °C which allowed attaining full reaction conversion at shorter time (24 h).

Entry	Catalyst	t [h]	Conv [%]	Yield [%] <sup>[b]</sup>	d.r.	ee [%] <sup>[c]</sup>
1	<b>C1</b>	72	100	72	70:30	30
2	<b>C2</b>	72	65	49	75:25	16
3	<b>C3</b>	72	70	46	80:20	68
4	<b>C4</b>	72	23	n.d.	n.d.	n.d.
5	<b>C5</b>	40	100	75	75:25	-12
6 <sup>[d]</sup>	<b>C6</b>	96	n.d.	40	>99:1	99
7	<b>C6</b>	24	100	81	>99:1	98

[a] Reactions conducted on a 0.2 mmol scale in 0.4 mL 1,2-DCE at 50 °C (molar ratio of **1/2a**/catalyst 1:1.5:0.1). [b] Yields of isolated product after column chromatography. [c] Determined by chiral HPLC analysis. [d] Reaction carried out at room temperature. n.d.: not determined.

Under these conditions (10 mol% **C6** in 1,2-DCE at 50 °C) the reaction of **1** worked equally well with an array of 2-aryl  $\alpha$ -cyanoacetates **2** to afford the corresponding adducts **5** as essentially single diastereomer in yields within the range from 62% to 95% and ee values greater than 95% in most cases. As Table 2 shows, these results seem to be independent upon the meta/para substitution pattern of the aromatic ring or their electron donating/withdrawing character. Entry 6 was an exception probably because of steric constraints imposed by the ortho substituent. Using the less sterically demanding benzyl and ethyl cyano esters **3** and **4**, a slight loss of stereoselection was produced (entries 9 and 10) albeit it was still acceptable. Configuration of adduct **5e** was established by a single crystal X-ray analysis<sup>[21]</sup> and that of the remaining adducts by assuming a uniform reaction mechanism.

**Double asymmetric induction.** Given the paucity of methods for the construction of tertiary/quaternary nonadjacent stereocenters, this organocatalytic Michael-protonation cascade was next extended to chiral  $\alpha'$ -oxy enones **8/9** and **10/11**,<sup>[22]</sup> a type of substrates that, as far as we know, have neither been previously investigated within the realm of organocatalysis.<sup>[23,24]</sup> As the results in Table 3 show, the stereochemical outcome of the reactions varies notably depending on the catalysts used. With Et<sub>3</sub>N as the only promoter (entries 1 and 8), adducts **12** and **14** were produced with moderate diastereoselectivity (ratio of *RSS*/other isomers 60:40 and 71:29, respectively). Again, the best catalyst was **C6** which afforded only two out of the four possible diastereomers in all the cases studied (entries 2, 3, 6, 9 and 10), with remarkable diastereoselectivity (*RSS*/*RRS*/others up to 91:9:0). Catalyst **C7** led to no reaction (entry 4), and **C8** and **C9** led to poor selectivity (entries 5 and 7). It is worth noting that in the above reactions *O*-silylated enones **10/11** behaved

**Table 2.** Scope of the conjugate addition of  $\alpha$ -cyanoacetates to **1**.<sup>[a]</sup>



Entry	R <sup>1</sup>	Product	t [h]	Yield [%] <sup>[b]</sup>	d.r.	ee [%] <sup>[c]</sup>
1		<b>5b</b>	24	69	98:2	98
2		<b>5c</b>	24	95	98:2	96
3		<b>5d</b>	40	70	99:1	>98
4		<b>5e</b>	40	67	98:2	>98
5		<b>5f</b>	40	83	>98:2	97
6		<b>5g</b>	40	NR	--	--
7		<b>5h</b>	16	62	97:3	96
8		<b>5i</b>	20	72	96:4	97
9		<b>6e</b>	24	76	90:10	92
10		<b>7b</b>	24	88	88:12	91

[a] Reactions conducted on a 0.2 mmol scale in 0.4 mL 1,2-DCE at 50 °C (molar ratio of **1**/cyanoester/**C6** 1:1.5:0.1). [b] Yields of isolated product after column chromatography. [c] Determined by chiral HPLC analysis.

superior than hydroxy enones **8/9** as the results in entries 2/3 (dr of 67:32 and 89:11, respectively) and 8/9 (dr of 71:29 and 91:9, respectively) show.<sup>[25]</sup>


At this point it remained unclear whether the above substrate/catalyst combinations correspond to a matched stereochemical relationship. To answer that question, the reaction between **2a** and the (*S*)-configured ent-**10** was carried out in the presence of catalyst **C6**. As the data in Scheme 3 show, a 69:12:12:7 mixture of diastereomers was obtained, with (*S,S,S*)-**16** as the major product. By comparison with data in entry 3 of Table 3, it seems clear that the pair **10/C6**, with the configurations (*R*)-substrate/(*S,S*)-catalyst, corresponds to the matched combination. Experiments using bifunctional Brønsted base catalysts derived from other chiral 1,2-primary/tertiary diamines also revealed (*R*)-substrate/(*S,S*)-catalyst as the best combination to induce formation of adducts of *RSS* configuration.<sup>[26]</sup>

**Table 3.** Michael addition/protonation cascade involving chiral enones (double asymmetric induction).<sup>[a]</sup>




**Scheme 3.** Reaction involving substrate/catalyst mismatched combination. TMS: trimethylsilyl.

Given these observations and the absence of studies concerning double asymmetric induction in this field, we next examined briefly the reaction of chiral α<sup>1</sup>-hydroxy enones without substituents at C<sub>α</sub> position. Under optimized conditions it was found that reaction of **17** with either **2b** or **2c** produced the corresponding adducts **18** and **19**, respectively, essentially as single diastereomers (Scheme 4). This result thus confirms that generation of the quaternary stereocenter proceeds with almost perfect asymmetric induction with catalyst **C6** for both α-substituted and unsubstituted enones.



**Scheme 4.** Generation of a quaternary γ-stereocenter in chiral α-unsubstituted α<sup>1</sup>-hydroxy ketones. TES: triethylsilyl.




**Scheme 5.** Elaboration of adducts: a) conversion of ketol into carboxy and aldehyde functions; b) stereoselective 1,2-diol formation.

**Adduct elaboration and proposed reaction models.** The results obtained are of special interest in that treatment of adducts **5a** and **5c** with NaIO<sub>4</sub> in MeOH/H<sub>2</sub>O provides the carboxylic acids **20** and **21** in 86% and 88% yield, respectively, along with acetone as the only organic side product formed (Scheme 5). Acid **21** was transformed into its methyl ester **22** for comparative purposes, *vide infra*. Alternatively, reduction of the carbonyl group of **5a** followed by diol cleavage as above furnished the aldehyde **23** in 76% yield over the two steps. Thus, the lack of reactivity and selectivity associated with methacrylate esters and methacrolein (*vide supra*) may now be remediated with this new methacrylate equivalent. Finally, to confirm the stereochemical assignment of reactions involving double asymmetric induction (Table 3), adduct **15** (90:10 diastereomeric mixture) was subjected to the oxidative

cleavage conditions to afford the same product **22** to that obtained from **5c**, along with the minor isomer **24**. Similarly, **18** upon oxidative cleavage of the ketol moiety as above and subsequent esterification of the resulting carboxylic acid provided the methyl ester **25**. The absolute configuration of both **24** and **25** was established by chemical correlation.<sup>[26]</sup> In addition to the above transformations, stereoarrays bearing up to four stereogenic centers may also be produced from this approach. Thus, diols **26** and **27** were obtained as essentially single *anti*-diol isomer through reduction of the respective  $\alpha'$ -hydroxy ketone **13** and **18** with  $\text{Zn}(\text{BH}_4)_2$ .<sup>[27]</sup>

The high fidelity with which chirality is transferred from the catalyst to the reaction products could be explained by the stereomodels depicted in Figure 4. By analogy to previously calculated TS geometries for the related conjugate addition of cyanoesters to  $\alpha$ -unsubstituted enone analog to **1**,<sup>[10]</sup> ternary complex **A** would account for the conjugate addition step, which would proceed with the catalyst interacting with both reaction components through several H-bonds. Once the addition adduct is formed, the local negative charge would no longer be located in the cyano ester moiety but in the enolate site. This will weaken the H-bond between the protonated quinuclidine and the cyano ester carbonyl. Finally, proton transfer, either directly from the protonated catalyst to the enolate or alternatively mediated by some proton-shuttle mechanism, would preferentially occur through the enolate *Re* face, as depicted in proposed model **B**.



**Figure 4.** Proposed approaching models for the addition and protonation steps, respectively.

## Conclusions

Direct approaches to the construction of acyclic carbonyl compounds with nonadjacent all carbon tertiary/quaternary stereocenters that proceed with high diastereo- and enantioselectivity are lacking. Here an effective solution to this longstanding problem is reported based on a bifunctional Brønsted base-catalyzed Michael/ $\alpha$ -protonation cascade that involves 2,4-dimethyl-4-hydroxypenten-3-one as design methacrylate equivalent. Key feature of this template is the ability to act as either H-bond donor or/and acceptor, a distinguishing feature among known bidentate enoate equivalents employed in organocatalysis.<sup>[11]</sup> Control experiments with other elementary Michael acceptors lacking such ambivalent character led to

inferior reactivity and/or selectivity. This design element also demonstrated successful in Michael/protonation cascades involving chiral  $\alpha'$ -oxyenones. In this latter case, double asymmetric induction occurs with substrate/catalyst matched combination providing adducts in up to >98:2 dr. The obtained adducts are easily transformed into the corresponding acyclic carboxylic acids, aldehydes and 1,2-diols with up to four configurationally-defined stereocenters. We believe this new family of enoate equivalents will rapidly find further applications in organocatalysis.

## Experimental Section

Selected experimental procedures. (All experimental details can be found in the Supporting Information. The material includes compound characterization, stereochemical determinations and copies of spectra of new compounds.)

**4-Hydroxy-2,4-dimethylpent-1-en-3-one (1):** To a solution of commercial methyl 2-hydroxy-2-methylpropanoate (15 mmol, 1.77 g) and *N,O*-dimethylhydroxylamine hydrochloride (22.5 mmol, 2.14 g, 1.5 equiv.) in THF (50 mL), a 2M solution of  $i\text{PrMgCl}$  in THF (60 mmol, 4 equiv.) was added at  $-20^\circ\text{C}$ . Once the reaction mixture was stirred for 1.5 h at room temperature, it was quenched with an aqueous saturated solution of  $\text{NH}_4\text{Cl}$  (30 mL) and extracted with  $\text{CH}_2\text{Cl}_2$  ( $2 \times 30$  mL). The combined organic phases were dried over  $\text{MgSO}_4$ . After filtration the solvent was evaporated under reduced pressure and the crude material was purified by flash column chromatography (eluent hexane/ethyl acetate 80/20) to obtain the desired Weinreb amide product. Yield: 1.99 g (90 %), colorless oil. To a solution of this material (10 mmol, 1.85 g) in  $\text{Et}_2\text{O}$  (20 mL) at  $-20^\circ\text{C}$ , a solution of isopropenyl magnesium bromide (0.5 M in THF, 60 mL, 3 equiv.) was added, and the resulting mixture was stirred at  $0^\circ\text{C}$  for 16 h. The reaction was quenched with an aqueous saturated solution of  $\text{NH}_4\text{Cl}$  (50 mL) and extracted with  $\text{Et}_2\text{O}$  ( $2 \times 50$  mL). The combined organic phases were dried over  $\text{MgSO}_4$ , filtered and the solvent was evaporated under reduced pressure. The crude product was purified by flash column chromatography (pentane/ $\text{Et}_2\text{O}$  95/5) to obtain the title compound. Yield: 833 mg (65 %), colorless oil.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were identical to those reported in the literature.<sup>[28]</sup>

**Preparation of chiral  $\alpha'$ -hydroxy enones 8/9:** To a solution of methyl 2-hydroxy-3-phenylpropanoate or methyl 2-hydroxy-4-methylpentanoate (10 mmol) and *N,O*-dimethylhydroxylamine hydrochloride (15 mmol, 1.5 equiv.) in THF (35 mL), at  $-20^\circ\text{C}$  a 2M solution of  $i\text{PrMgCl}$  in THF (40 mmol, 20 mL, 4 equiv.) was added. The reaction mixture was stirred for 1.5 h at  $0^\circ\text{C}$ . The reaction was then quenched with an aqueous saturated solution of  $\text{NH}_4\text{Cl}$  (30 mL) and extracted with  $\text{CH}_2\text{Cl}_2$  ( $2 \times 30$  mL). The combined organic phases were dried over  $\text{MgSO}_4$ . After filtration the solvent was evaporated under reduced pressure and the crude product was purified by flash column chromatography (eluent hexane/ethyl acetate 80/20) to obtain the corresponding Weinreb amide.

To a solution of 2-bromopropene (9 mmol, 0.79 mL, 3 equiv.) in  $\text{Et}_2\text{O}$  (5 mL) at  $-78^\circ\text{C}$ , a solution of *tert*-butyllithium (1.6M in pentane, 6.75 mL, 3.6 equiv.) was added, and the resulting mixture was stirred at the same temperature for 1 h. Subsequently, a solution of the corresponding Weinreb amide (3 mmol) in  $\text{Et}_2\text{O}$  (10 mL) was added and the reaction mixture was stirred at  $-60^\circ\text{C}$  for 16 h. The reaction was quenched with an aqueous saturated solution of  $\text{NH}_4\text{Cl}$  (50 mL) and extracted with  $\text{CH}_2\text{Cl}_2$  (50 mL). The organic phase was dried over  $\text{MgSO}_4$ , filtered and the solvent was evaporated under reduced pressure. The crude product was purified by flash column chromatography (eluent hexane/ethyl acetate 95/5).

**Preparation of chiral  $\alpha$ '-silyloxy enones 10/11:** To a solution of the corresponding  $\alpha$ '-hydroxy enone **8/9** (2 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 mL) at  $-20^\circ\text{C}$ , were added successively 2,6-lutidine (0.55 mL, 4.8 mmol, 2.4 equiv.) and TMSOTf (0.72 mL, 4 mmol, 2 equiv.), and the mixture was stirred at the same temperature for 3 h. EtOAc (40 mL) was then added, and the organic phase was washed with saturated aqueous solutions of  $\text{NaHCO}_3$  (40 mL),  $\text{CuSO}_4$  ( $3 \times 40$  mL),  $\text{NaHCO}_3$  ( $2 \times 40$  mL) and  $\text{NaCl}$  (40 mL). The organic phase was dried over  $\text{MgSO}_4$ , filtered and the solvent was evaporated under reduced pressure. The resulting crude material was purified by flash column chromatography (eluent hexane/ethyl acetate 99/1) to obtain pure compounds **10/11**.

**General Procedure for the catalytic conjugate addition of  $\alpha$ -cyanoacetates to **1**:** To a solution of the corresponding  $\alpha$ -cyanoacetate **2** (0.3 mmol, 1.5 equiv.) and  $\alpha$ '-hydroxy enone **1** (26 mg, 0.2 mmol) in 1,2-DCE (0.4 mL), catalyst **C6** (13 mg, 0.02 mmol) was added, and the resulting mixture was stirred at  $50^\circ\text{C}$  until consumption of enone **1** (monitored by  $^1\text{H-NMR}$ ). Then the reaction was quenched with  $\text{HCl}$  1N and the mixture was extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 2$  mL). The combined organic phases were dried over  $\text{MgSO}_4$  and the solvent was removed under reduced pressure to give the corresponding addition/ $\alpha$ -protonation adduct which was purified by flash column chromatography (eluent hexane/ethyl acetate 95/5).

**Catalytic conjugate addition of  $\alpha$ -cyanoacetates **2** to chiral  $\alpha$ '-oxy enones:** To a solution of the corresponding *tert*-butyl cyanoacetate **2** (0.6 mmol) and the corresponding  $\alpha$ '-oxy enone **8-11** (0.2 mmol, 1 equiv.) in  $\text{CH}_2\text{Cl}_2$  (0.4 mL), the catalyst (0.02 mmol) was added and the resulting mixture was stirred at  $20^\circ\text{C}$  until consumption of the  $\alpha$ '-oxy enone (monitored by  $^1\text{H-NMR}$ ; see Table 3 for reaction times). The reaction mixture was quenched with  $\text{HCl}$  1N (5 mL) and the solution was extracted with  $\text{CH}_2\text{Cl}_2$  (5 mL). The combined organic phases were dried over  $\text{MgSO}_4$ , filtered and the solvent was evaporated under reduced pressure.

Reactions from  $\alpha$ '-hydroxy enone **8/9**: The residue was submitted to purification by flash column chromatography (eluent hexane/ethyl acetate 95/5).

Reactions from  $\alpha$ '-silyloxy enone **10/11**: The residue was dissolved in MeOH (0.5 mL) and a solution of concentrated fluorhydric acid in MeOH (10 mmol, 0.2 mL) was added. The resulting mixture was stirred at  $20^\circ\text{C}$  for 2 h. Then the solvent was evaporated and the resulting residue was basified to pH 7 with a saturated solution of  $\text{NaHCO}_3$ . The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  ( $2 \times 4$  mL), dried over  $\text{MgSO}_4$ , filtered and the solvent was evaporated under reduced pressure. The oily product was submitted to purification by flash column chromatography (eluent hexane/ethyl acetate 95/5).

**Conversion of adduct **5a** into carboxylic acid **20**:** A suspension of sodium periodate  $\text{NaIO}_4$  (342 mg, 1.6 mmol) in water (0.8 mL) was added to a solution of  $\alpha$ -hydroxy ketone **5a** (69 mg, 0.2 mmol) in methanol (1 mL). The mixture was stirred at room temperature until the reaction was complete (monitored by TLC, 24 h). Then the solvent was removed under reduced pressure. Water (4.5 mL) was added to the residue and the resulting mixture was extracted with  $\text{Et}_2\text{O}$  ( $3 \times 6$  mL). The combined organic extracts were dried over  $\text{MgSO}_4$ , filtered and the solvent was evaporated. The product was purified by flash column chromatography (Hex:EtOAc 80:20) to afford carboxylic acid **20** (52 mg, 86% yield, colorless oil).

**Conversion of adduct **5a** into aldehyde **23**:**  $\text{BH}_3\cdot\text{THF}$  (1 M, 0.4 mL, 0.4 mmol) was added to a solution of  $\alpha$ -hydroxy ketone **5a** (69 mg, 0.2 mmol) in dry THF (0.9 mL) at  $0^\circ\text{C}$  and the resulting solution was stirred at the same temperature for 2 h. Then MeOH (1 mL) was added and the resulting mixture was stirred at room temperature for 30 min. The solvents were removed under reduced pressure and the residue thus obtained was subjected to oxidative scission by treatment with  $\text{NaIO}_4$ , as above. The crude material was purified by flash column chromatography on silica gel (eluting with Hex:EtOAc 95/5) to give the title compound (44 mg, 76% yield, oil).

## Acknowledgements

Support was provided by the University of the Basque Country UPV/EHU (UFI 11/22), Gobierno Vasco (GV, Grant No IT-628-13), and Ministerio de Economía y Competitividad (MEC, Grant CTQ2013-47925-C2), Spain. E. B. thanks MEC and I. I. thanks GV for Fellowships. We also thank SGIker (UPV/EHU) for providing NMR, HRMS, and X-Ray resources.

**Keywords:** asymmetric organocatalysis • Michael reactions • stereoselective protonation • quaternary stereocenters • Brønsted bases

- [1] a) K. L. Kimmel, J. D. Weaver, M. Lee, J. A. Ellman, *J. Am. Chem. Soc.* **2012**, *134*, 9058-9061; b) B. Zheng, H. Wang, Y. Han, C. Liu, Y. Peng, *Chem. Commun.* **2013**, *49*, 4561-4563; c) N. V. G. Moorthy, R. Dyapa, S. V. Pansare, *Org. Lett.* **2015**, *17*, 5312-5315.
- [2] For classic examples of multistep approaches to  $\alpha,\gamma$ -substituted carbonyl patterns en route to erythromycins, see: a) E. J. Corey, P. B. Hopkins, S. K. Sung-eun, Y. Krishnan, P. Nambiar, J. R. Falck, *J. Am. Chem. Soc.* **1979**, *101*, 7131-7134; b) R. B. Woodward *et al.*, *J. Am. Chem. Soc.* **1981**, *103*, 3215-3217; c) G. Stork, D. R. Rychnovsky, *J. Am. Chem. Soc.* **1987**, *109*, 1565-1567; d) J. Mulzer, *Angew. Chem.* **1991**, *103*, 1484-1486; *Angew. Chem. Int. Ed.* **1991**, *30*, 1452-1454; e) R. Stürmer, K. Ritter, R. W. Hoffmann, *Angew. Chem.* **1993**, *105*, 112-114; *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 101-103.
- [3] For a racemic synthesis of a  $\alpha,\gamma$ -substituted carbonyl pattern with all-carbon quaternary/tertiary nonadjacent stereocenters, see: J.-H. Fan, W.-T. Wei, M.-B. Zhou, R.-J. Song, J.-H. Li, *Angew. Chem.* **2014**, *124*, 6768-6772; *Angew. Chem. Int. Ed.* **2014**, *53*, 6650-6654.
- [4] a) Y. Wang, X. Liu, L. Deng, *J. Am. Chem. Soc.* **2006**, *128*, 3928-3930; b) B. Wang, F. Wu, Y. Wang, X. Liu, L. Deng, *J. Am. Chem. Soc.* **2007**, *129*, 768-769. For an extension to 2-oxindoles as pronucleophiles, see: c) X. Li, S. Luo, J.-P. Cheng, *Chem. Eur. J.* **2010**, *16*, 14290-14294.
- [5] a) S.-W. Duan, J. An, J. R. Chen, W. J. Xiao, *Org. Lett.* **2011**, *13*, 2290-2293; b) S.-W. Duan, Y.-Y. Liu, W. Ding, T.-R. Li, D.-Q. Shi, J. R. Chen, W. J. Xiao, *Synthesis* **2013**, *45*, 1647-1653.
- [6] T. Tsubogo, S. Saito, K. Seki, Y. Yamashita, S. Kobayashi, *J. Am. Chem. Soc.* **2008**, *130*, 13321-13332.
- [7] a) E. K. Kemppainen, G. Sahoo, A. Valkonen, P. M. Pihko, *Org. Lett.* **2012**, *14*, 1086-1089; b) E. K. Kemppainen, G. Sahoo, A. Piisola, A. Hamza, B. Kótai, I. Pápai, P. M. Pihko, *Chem. Eur. J.* **2014**, *20*, 5983-5993. Also, see: c) N. Fu, L. Zhang, S. Luo, J.-P. Cheng, *Chem. Eur. J.* **2013**, *19*, 15669-15681.
- [8] For a recent approach to overcome problems associated with the enantioselective conjugate addition to methacrylate esters, see: a) A. J. M. Farley, C. Sandford, D. J. Dixon, *J. Am. Chem. Soc.* **2015**, *137*, 15992-15995. For recent reviews on conjugate addition/enantioselective protonation, see: b) S. Oudeyer, J.-F. Brière, V. Levacher, *Eur. J. Org. Chem.* **2014**, 6103-6119; c) J. T. Mohr, A. Y. Hong, B. M. Stoltz, *Nat. Chem.* **2009**, *1*, 359-369; d) J. P. Phelan, J. A. Ellman, *Belstein J. Org. Chem.* **2016**, *12*, 1203-1228.
- [9] For the interest in  $\alpha$ -functionalized 2-cyanoacetates, see: a) S. Jautza, R. Peters, *Synthesis* **2010**, 365-388; b) M. D. Díaz de Villegas, J. A. Gálvez, R. Badorrey, P. López Ram de Viv, *Adv. Synth. Catal.* **2014**, *356*, 3261-3288.
- [10] E. Badiola, B. Fiser, E. Gómez-Bengoia, A. Mielgo, I. Olaizola, I. Urruzuno, J. M. García, J. M. Odriozola, J. Razkin, M. Oiarbide, C. Palomo, *J. Am. Chem. Soc.* **2014**, *136*, 17869-17881.
- [11] Recent review on unsaturated carboxylic acid equivalents in asymmetric organocatalysis: a) D. Monge, H. Jiang, Y. Alvarez-Casao, *Chem. Eur.*

- J.* **2015**, *21*, 4494-4504. Use of  $\alpha,\beta$ -unsaturated amides and related systems in conjugate additions: b) K. M. Byrd, *Belstein J. Org. Chem.* **2015**, *11*, 530-562; c) G. Desimoni, G. Faita, P. Quadrelli, *Chem. Rev.* **2015**, *115*, 9922-9980.
- [12] Unsaturated carbonyl compounds used as Michael acceptors in conjugate addition/enantioselective protonation usually are cyclic, because the cycle unequivocally determines the configuration of evolving enolate. For some examples of acyclic acceptors leading to a single stereocenter: (iminium activation with enals) a) N. Fu, L. Zhang, J. Li, S. Luo, J.-P. Cheng, *Angew. Chem.* **2011**, *123*, 10061-10063; *Angew. Chem. Int. Ed.* **2011**, *50*, 11451-11455; (iminium activation with enones) b) N. Fu, L. Zhang, S. Luo, J.-P. Cheng, *Chem. Eur. J.* **2013**, *19*, 15669-15681; c) L. Cui, L. Zhang, S. Luo, J.-P. Cheng, *Eur. J. Org. Chem.* **2014**, 3540-3545; d) N. Fu, L. Zhang, S. Luo, *Org. Lett.* **2015**, *17*, 382-385; (Zn(II) or Ca(II) catalysis with unsaturated imides) e) M. P. Sibi, J. Coulomb, L. M. Stanley, *Angew. Chem.* **2008**, *120*, 10061-10063; *Angew. Chem. Int. Ed.* **2008**, *47*, 9913-9915; f) T. Poisson, Y. Yamashita, S. Kobayashi, *J. Am. Chem. Soc.* **2010**, *132*, 7890-7892. (Rh catalysis with dehydro amino acids) g) L. Navarre, R. Martinez, J.-P. Genet, S. Darses, *J. Am. Chem. Soc.* **2008**, *130*, 6159-6169. Acyclic acceptor leading to two contiguous stereocenters: h) M. P. Sibi, G. Petrovic, J. Zimmerman, *J. Am. Chem. Soc.* **2005**, *127*, 2390-2391. For further examples, see reference 8.
- [13] During the preparation of this manuscript, an organocatalytic conjugate addition/protonation sequence leading to nonadjacent stereocenters was reported, although using a cyclic Michael acceptor (N-itaconimide): B. Zhu, R. Lee, J. Li, X. Ye, S.-N. Hong, S. Qiu, M. L. Coote, Z. Jiang, *Angew. Chem.* **2016**, *128*, 1321-1325; *Angew. Chem. Int. Ed.* **2016**, *55*, 1299-1303.
- [14] Use of  $\alpha$ -hydroxy enones in asymmetric transformations: C. Palomo, M. Oiarbide, J. M. García, *Chem. Soc. Rev.* **2012**, *41*, 4150-4164.
- [15] This enone could be prepared from commercial methyl 2-hydroxy-2-methylpropanoate and isopropenylmagnesium bromide in two steps (59% overall yield). See the Supporting Information for details.
- [16] Reviews on Brønsted bases: a) C. Palomo, M. Oiarbide, R. López, *Chem. Soc. Rev.* **2009**, *38*, 632-653; b) *Science of Synthesis: Asymmetric Organocatalysis 2, Brønsted Base and Acid Catalysis, and Additional Topics* (Ed.: K. Maruoka), Georg Thieme, Stuttgart, **2012**; c) A. Ting, J. M. Gross, N. T. McDougal and S. E. Schaus, *Top. Curr. Chem.*, **2010**, *291*, 145-200.
- [17] a) S. H. McCooley, S. Connon, *Angew. Chem.* **2005**, *117*, 6525-6528; *Angew. Chem. Int. Ed.* **2005**, *44*, 6367-6370; b) J. Ye, D. J. Dixon, P. S. Hynes, *Chem. Commun.* **2005**, 4481-4483; c) B. Vakulya, S. Varga, A. Csampai, T. Soós, *Org. Lett.* **2005**, *7*, 1967-1969; d) B.-J. Li, L. Jiang, M. Liu, Y.-C. Chen, L.-S. Ding, Y. Wu, *Synlett* **2005**, 603-606.
- [18] K. Hu, A. Lu, Y. Wang, Z. Zhou, C. Tang, *Tetrahedron: Asymmetry* **2013**, *24*, 953-957.
- [19] a) J. P. Malerich, K. Hagihara, V. R. Rawal, *J. Am. Chem. Soc.* **2008**, *130*, 14416-14417; b) Y. Zhu, J. P. Malerich, V. H. Rawal, *Angew. Chem.* **2010**, *122*, 157-160; *Angew. Chem. Int. Ed.* **2010**, *49*, 153-156.
- [20] a) L. Dai, S.-X. Wang, F.-E. Chen, *Adv. Synth. Catal.* **2010**, *352*, 2137-2141; b) W. Yang, D.-M. Du, *Org. Lett.* **2010**, *12*, 5450-5453.
- [21] CCDC-1470018 (compound **5e**) contains the supplementary crystallographic data for this paper. These data can be obtained from the Cambridge Crystallographic Data Center via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif). See the Supporting Information for details.
- [22] These enones were easily prepared from the corresponding  $\alpha$ -hydroxy acids. See the Supporting Information for details.
- [23] *Science of Synthesis: Asymmetric Organocatalysis, Vol. 1 & 2* (Eds.: B. List, K. Maruoka), Georg Thieme, Stuttgart, **2012**.
- [24] Reactions with chiral  $\alpha$ -unsubstituted  $\alpha'$ -hydroxy enones leading to tertiary stereocenters: (Conjugate additions) a) W.-Y. Zhang, D. J. Jakiela, A. Maul, C. Knors, J. W. Lauher, P. Helquist, D. Enders, *J. Am. Chem. Soc.* **1988**, *110*, 4652-4660; b) E. J. Corey, F. J. Hannon, N. W. Boaz, *Tetrahedron* **1989**, *45*, 545-555; c) J. M. García, M. A. Maestro, M. Oiarbide, J. M. Odriozola, J. Razkin, C. Palomo, *Org. Lett.* **2009**, *11*, 3826-3829; d) L. C. Kopel, M. S. Ahmed, F. T. Halaweish, *Steroids* **2013**, *78*, 1119-1125; (Diels-Alder) e) W. Choy, L. A. Reed, III, S. Masamune, *J. Org. Chem.* **1983**, *48*, 1137-1139; f) S. Masamune, L. A. Reed, III, J. T. Davis, W. Choy, *J. Org. Chem.* **1983**, *48*, 4441-4444; g) B. Stammen, U. Berlage, R. Kinderman, M. Kaiser, B. Günther, W. S. Sheldrick, P. Welzel, W. R. Roth, *J. Org. Chem.* **1992**, *57*, 6566-6575; h) P. Bañuelos, J. M. García, E. Gómez-Bengoia, A. Herrero, J. M. Odriozola, M. Oiarbide, C. Palomo, J. Razkin, *J. Org. Chem.* **2010**, *75*, 1458-1473. (Cycloaddition with nitrones) i) C. Palomo, M. Oiarbide, E. Arceo, J. M. García, R. López, A. González, A. Linden, *Angew. Chem.* **2005**, *117*, 6343-6346; *Angew. Chem. Int. Ed.* **2005**, *44*, 6187-6190.
- [25] No racemization occurred when **8** and **10** were stirred in the presence of Et<sub>3</sub>N, **C6** or DBU. Partial epimerization of product **13** appeared only upon prolonged exposure to DBU. See the Supporting Information for details.
- [26] See the Supporting Information for details.
- [27] T. Nakata, T. Tanaka, T. Oishi, *Tetrahedron Lett.* **1983**, *24*, 2653-2656.
- [28] A. Basheer, M. Mishima, I. Marek, *Org. Lett.* **2011**, *13*, 4076-4079.

## FULL PAPER

## Entry for the Table of Contents (Please choose one layout)

Layout 1:

## FULL PAPER

Text for Table of Contents

Author(s), Corresponding Author(s)\*


Page No. – Page No.

Title

((Insert TOC Graphic here: max. width: 5.5 cm; max. height: 5.0 cm))

Layout 2:

## FULL PAPER



2,4-Dimethyl-4-hydroxypenten-3-one is introduced as new methacrylate equivalent for the asymmetric construction of nonadjacent all carbon tertiary/quaternary stereocenters via organocatalytic addition of  $\alpha$ -cyanoacetates. Direct assembly of such stereoarrays with high enantio- and diastereoselectivity has remained elusive to date.

Igor Iriarte, Silvia Vera, Eider Badiola, Antonia Mielgo, Mikel Oiarbide,\* Jesús M. García, José M. Odriozola, and Claudio Palomo\*

**Page No. – Page No.**  
**Asymmetric Assembly of All-Carbon Tertiary/Quaternary Nonadjacent Stereocenters Through Organocatalytic Conjugate Addition of  $\alpha$ -Cyanoacetates to a Methacrylate Equivalent.**