

## eReferences

- e1. Friedrich J, Sorge S, Bujupi F, et al. Hox Function Is Required for the Development and Maintenance of the Drosophila Feeding Motor Unit. *Cell Rep.* Feb 02 2016;14(4):850-860. doi:10.1016/j.celrep.2015.12.077
- e2. Bacalini MG, Gentilini D, Boattini A, et al. Identification of a DNA methylation signature in blood cells from persons with Down Syndrome. *Aging (Albany NY).* Feb 2015;7(2):82-96. doi:10.18632/aging.100715
- e3. Fuso A, Ferraguti G, Grandoni F, et al. Early demethylation of non-CpG, CpC-rich, elements in the myogenin 5'-flanking region: a priming effect on the spreading of active demethylation. *Cell Cycle.* Oct 01 2010;9(19):3965-76. doi:10.4161/cc.9.19.13193
- e4. Monti N, Cavallaro RA, Stoccoro A, et al. CpG and non-CpG Presenilin1 methylation pattern in course of neurodevelopment and neurodegeneration is associated with gene expression in human and murine brain. *Epigenetics.* 08 2020;15(8):781-799. doi:10.1080/15592294.2020.1722917
- e5. Nicolai V, Ciraci V, Cavallaro RA, Ferrer I, Scarpa S, Fuso A. GSK3 $\beta$  5'-flanking DNA Methylation and Expression in Alzheimer's Disease Patients. *Curr Alzheimer Res.* 2017;14(7):753-759. doi:10.2174/1567205014666170203153325
- e6. Fedotova EY, Iakovenko EV, Abramychева NY, Illarioshkin SN. Gene Methylation in Parkinson's Disease and Multiple System Atrophy. *Epigenomes.* Feb 06 2023;7(1)doi:10.3390/epigenomes7010005
- e7. Mercurio R, Pergoli L, Galimberti D, et al. PICALM Gene Methylation in Blood of Alzheimer's Disease Patients Is Associated with Cognitive Decline. *J Alzheimers Dis.* 2018;65(1):283-292. doi:10.3233/JAD-180242
- e8. Nicolai V, Cavallaro RA, López-González I, et al. DNA Methylation Profiles of Selected Pro-Inflammatory Cytokines in Alzheimer Disease. *J Neuropathol Exp Neurol.* Jan 01 2017;76(1):27-31. doi:10.1093/jnen/nlw099